

KLHL7 Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP9307c

Specification

KLHL7 Antibody (Center) Blocking Peptide - Product Information

Primary Accession

08IXQ5

KLHL7 Antibody (Center) Blocking Peptide - Additional Information

Gene ID 55975

Other Names

Kelch-like protein 7, KLHL7

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP9307c was selected from the Center region of human KLHL7. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

KLHL7 Antibody (Center) Blocking Peptide - Protein Information

Name KLHL7

Function

Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin ligase complex. The BCR(KLHL7) complex acts by mediating ubiquitination and subsequent degradation of substrate proteins. Probably mediates 'Lys-48'-linked ubiquitination.

Cellular Location

Nucleus. Cytoplasm. Note=Colocalizes with CUL3 in punctate structures at the perinuclear region of the cytoplasm

Tissue Location

Widely expressed, with highest levels in adult and fetal heart, CNS and adult testis.



Tel: 858.875.1900 Fax: 858.875.1999

KLHL7 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

KLHL7 Antibody (Center) Blocking Peptide - Images

KLHL7 Antibody (Center) Blocking Peptide - Background

KLHL7 encodes a BTB-Kelch-related protein. The encoded protein may be involved in protein degradation.

KLHL7 Antibody (Center) Blocking Peptide - References

Friedman, J.S., et.al., Am. J. Hum. Genet. 84 (6), 792-800 (2009) Bredholt, G., et.al, Scand. J. Immunol. 64 (3), 325-335 (2006)