10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CBAA1 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP9682a

Specification

CBAA1 Antibody (N-term) Blocking Peptide - Product Information

Primary Accession Q9BPX6

CBAA1 Antibody (N-term) Blocking Peptide - Additional Information

Gene ID 10367

Other Names
Calcium uptake protein 1, mitochondrial, Atopy-related autoantigen CALC, ara CALC, Calcium-binding atopy-related autoantigen 1, Hom s 4, MICU1, CALC, CBARA1

Format
Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions
This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CBAA1 Antibody (N-term) Blocking Peptide - Protein Information

Name MICU1 {ECO:0000303|PubMed:20693986, ECO:0000312|HGNC:HGNC:1530}

Function
Calcium sensor of the mitochondrial calcium uniporter (MCU) channel, which senses calcium level via its EF-hand domains (PubMed:20693986, PubMed:23101630, PubMed:23747253, PubMed:24313810, PubMed:24332854, PubMed:24503055, PubMed:24560927, PubMed:26341627, PubMed:26903221, PubMed:27099988, PubMed:28615291, PubMed:30454562, PubMed:30638448, PubMed:32494073, PubMed:32667285, PubMed:<a href="http://www.uniprot.org/citations/32762847"

target="_blank">32762847, PubMed:32790952, PubMed:34463251, PubMed:36206740, PubMed:37036971, PubMed:37126688). MICU1 and MICU2 (or MICU3) form a disulfide-linked heterodimer that stimulates and inhibits MCU activity, depending on the concentration of calcium (PubMed:24560927, PubMed:26903221, PubMed:28615291, PubMed:32148862, PubMed:32494073, PubMed:32667285, PubMed:32762847, PubMed:32790952, PubMed:36206740, PubMed:37036971, PubMed:37126688). At low calcium levels, MICU1 occludes the pore of the MCU channel, preventing mitochondrial calcium uptake (PubMed:32494073, PubMed:32667285, PubMed:32762847, PubMed:37036971, PubMed:37126688). At higher calcium levels, calcium-binding to MICU1 and MICU2 (or MICU3) induces a conformational change that weakens MCU-MICU1 interactions and moves the MICU1-MICU2 heterodimer away from the pore, allowing calcium permeation through the MCU channel (PubMed:32494073, PubMed:32667285, PubMed:32762847). Also required to protect against manganese toxicity by preventing manganese uptake by MCU: mechanistically, manganese- binding to its EF-hand domains does not induce any conformational change, maintaining MCU pore occlusion (PubMed:30082385, PubMed:30403999). Also acts as a barrier for inhibitors of the MCU channel, such as ruthenium red or its derivative Ru360 (PubMed:37244260). Acts as a regulator of mitochondrial cristae structure independently of its ability to regulate the mitochondrial calcium uniporter channel (PubMed:31427612, PubMed:37098122). Regulates glucose-dependent insulin secretion in pancreatic beta-cells by regulating mitochondrial calcium uptake (PubMed:22904319). Induces T- helper 1-mediated autoreactivity, which is accompanied by the release of IFNG (PubMed:16002733).

Cellular Location
Mitochondrion intermembrane space. Mitochondrion inner membrane. Note=Recruited to the mitochondrial inner membrane by EMRE/SMDT1 (PubMed:30454562). Also localizes to mitochondrial cristae junctions (PubMed:31427612)

Tissue Location
Expressed in epithelial cell lines. Strongly expressed in epidermal keratinocytes and dermal endothelial cells

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CBAA1 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Blocking Peptides

CBAA1 Antibody (N-term) Blocking Peptide - Images

CBAA1 Antibody (N-term) Blocking Peptide - References

Bordicchia, M., et al. Metab. Clin. Exp. (2009) In press :Grupe, A., et al. Am. J. Hum. Genet. 78(1):78-88(2006)Natter, S., et al. FASEB J. 12(14):1559-1569(1998)