

**ERC2 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP9944b****Specification**

---

**ERC2 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [O15083](#)**ERC2 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 26059**Other Names**

ERC protein 2, ERC2, KIAA0378

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**ERC2 Antibody (C-term) Blocking Peptide - Protein Information****Name** ERC2**Synonyms** KIAA0378**Function**

Thought to be involved in the organization of the cytomatrix at the nerve terminals active zone (CAZ) which regulates neurotransmitter release. Seems to act together with BSN. May recruit liprin-alpha proteins to the CAZ.

**Cellular Location**

Cytoplasm. Synapse. Presynaptic active zone. Cytoplasm, cytoskeleton. Note=In neurons, localized to synapses, and colocalizes with PCLO. Localized to the active zone of presynaptic density (By similarity).

**ERC2 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**ERC2 Antibody (C-term) Blocking Peptide - Images****ERC2 Antibody (C-term) Blocking Peptide - Background**

ERCs (ERC1 and ERC2) also referred to as ELKS and CAST are related proteins which share an identical C-terminal sequence. They interact with the conserved RIM PDZ domain via an unusual PDZ binding motif. ERC2/CAST 1 is only expressed as a single RIM binding variant. All ERCs have been shown to interact with Rab6, a protein involved in membrane trafficking at the Golgi complex. The function of these proteins has not been determined yet. They may link Rab6 mediated non-neuronal membrane traffic at the Golgi complex to neuronal membrane traffic at the active zone executed via RIMs.

**ERC2 Antibody (C-term) Blocking Peptide - References**

Uhl, G.R., et al. Arch. Gen. Psychiatry 65(6):683-693(2008)Kiel, D.P., et al. BMC Med. Genet. 8 SUPPL 1, S14 (2007) tom Dieck, S., et al. J. Cell Biol. 168(5):825-836(2005)