

Recombinant Human FGF-basic (154 a.a.)
Catalog # PBG10111**Specification**

Recombinant Human FGF-basic (154 a.a.) - Product Information**Recombinant Human FGF-basic (154 a.a.) - Additional Information****Description**

FGF-basic is one of 23 known members of the FGF family. Proteins of this family play a central role during prenatal development and postnatal growth and regeneration of a variety of tissues, by promoting cellular proliferation and differentiation. FGF-basic is a non-glycosylated heparin binding growth factor that is expressed in the brain, pituitary, kidney, retina, bone, testis, adrenal gland liver, monocytes, epithelial cells and endothelial cells. FGF-basic signals through FGFR 1b, 1c, 2c, 3c and 4. Recombinant human FGF-basic is a 17.2 kDa protein consisting of 154 amino acid residues.

BiologicalActivity

Assay #1: The **ED₅₀** as determined by the dose-dependent stimulation of thymidine uptake by BaF3 cells expressing FGF receptors is ≤ 0.5 ng/ml, corresponding to a specific activity of $\geq 2 \times 10^6$ units/mg.
Assay #2: The **ED₅₀** was determined by a cell proliferation assay using balb/c 3T3 cells is ≤ 0.1 ng/ml, corresponding to a specific activity of $\geq 1 \times 10^7$ units/mg.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is <0.1 ng/ μ g of protein (<1 EU/ μ g).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C

Precautions

Recombinant Human FGF-basic (154 a.a.) is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Human FGF-basic (154 a.a.) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Recombinant Human FGF-basic (154 a.a.) - Images