

Recombinant Human HCC-1 (CCL14) (66 a.a.)
Catalog # PBG10154**Specification**

Recombinant Human HCC-1 (CCL14) (66 a.a.) - Product Information**Recombinant Human HCC-1 (CCL14) (66 a.a.) - Additional Information****Description**

HCC-1 is a CC chemokine that signals through the CCR1 receptor and chemoattracts blood monocytes. It is secreted by various tissues including skeletal muscle, heart, spleen, liver, bone marrow and plasma. Mature HCC-1 is found in four different forms, which are distinguished by differential N-terminal truncation and contain 74, 72, 71, or 66 amino acid residues. Recombinant human HCC-1 (66 a.a.) is a 7.8 kDa protein consisting of 66 amino acids including the four highly conserved residues present in CC chemokines.

Biological Activity

Determined by its ability to chemoattract human monocytes using a concentration range of 5.0-20.0 ng/ml.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is <0.1 ng/ µg of protein (<1EU/ µg).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C

Precautions

Recombinant Human HCC-1 (CCL14) (66 a.a.) is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Human HCC-1 (CCL14) (66 a.a.) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Recombinant Human HCC-1 (CCL14) (66 a.a.) - Images