

Recombinant Human IGF-BP2
Catalog # PBG10171**Specification**

Recombinant Human IGF-BP2 - Product Information**Recombinant Human IGF-BP2 - Additional Information****Description**

IGF-BPs controls the distribution, function and activity of IGFs in various cell tissues and body fluids. Currently there are seven named IGF-BPs that form high affinity complexes with both IGF-I and IGF-II. IGF-BP2 is a cysteine-rich secreted protein produced by bone cells, and is most abundant in the brain. IGF-BP2 has been shown to inhibit IGF-II action in human breast and ovarian carcinoma cells. Recombinant human IGF-BP2 is a 31.5 kDa protein consisting of 289 amino acid residues including the IGF-BP domain and thyroglobulin type-I domain.
*Manufactured using (BTI-Tn-5B1-4) cells under license from the Boyce Thompson Institute for Plant Research, Inc.

BiologicalActivity

Determined by its ability to inhibit IGF-I induced proliferation of FDC-P1 cells.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is <0.1 ng/ µg of protein (<1EU/ µg).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C

Precautions

Recombinant Human IGF-BP2 is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Human IGF-BP2 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

Recombinant Human IGF-BP2 - Images