

Recombinant Human IL-16 (121 a.a.)
Catalog # PBG10197**Specification**

Recombinant Human IL-16 (121 a.a.) - Product Information**Recombinant Human IL-16 (121 a.a.) - Additional Information****Description**

IL-16 is a CD8+ T cell-derived cytokine that induces chemotaxis of CD4+ T cells and CD4+ monocytes and eosinophils. Analysis by gel filtration suggests that, under physiological conditions, hIL-16 exists predominantly as a noncovalently linked multimer, but that some IL-16 may exist as a monomer. However, only the multimeric form appears to possess chemotactic activity, suggesting that receptor cross-linking may be required for activity. IL-16 also induces expression of IL-2 receptor (IL-2R) and MHC class II molecules on CD4 + T cells. Human and murine IL-16 show significant cross-species reactivity. Recombinant human IL-16 is a 13.3 kDa protein consisting of 129 amino acid residues.

Biological Activity

Determined by its ability to chemoattract human CD4+ T-Lymphocytes using a concentration range of 50.0-100.0 ng/ml.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is <0.1 ng/ µg of protein (<1EU/ µg).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C

Precautions

Recombinant Human IL-16 (121 a.a.) is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Human IL-16 (121 a.a.) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

Recombinant Human IL-16 (121 a.a.) - Images