

Recombinant Murine IL-1a

Catalog # PBG10208

Specification

Recombinant Murine IL-1α - Product Information

Recombinant Murine IL-1α - Additional Information

Description

IL-1 α is a non-secreted proinflammatory cytokine produced in a variety of cells including monocytes, tissue macrophages, keratinocytes and other epithelial cells. Both IL-1 α and IL-1 β binds to the same receptor and has similar if not identical biological properties. These cytokines have a broad range of activities including, stimulation of thymocyte proliferation, by inducing IL-2 release, B-cell maturation and proliferation, mitogenic FGF-like activity and the ability to stimulate the release of prostaglandin and collagenase from synovial cells. However, whereas IL-1 β is a secreted cytokine, IL-1 α is predominantly a cell-associated cytokine. Recombinant murine IL-1 α is an 17.9 kDa protein containing 156 amino acid residues.

BiologicalActivity

The ED₅₀ as determined by the dose-dependent stimulation of murine D10S cells is < 0.002 ng/ml, corresponding to a specific activity of $> 5 \times 10 < \text{sup} > 8 < \text{sup} > \text{units/mg}$.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is $<0.1 \text{ ng}/\mu\text{g}$ of protein ($<1\text{EU}/\mu\text{g}$).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C

Precautions

Recombinant Murine IL- 1α is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Murine IL-1α - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation





- Flow CytometyCell Culture

Recombinant Murine IL- 1α - Images