

Recombinant Human NP-1
Catalog # PBG10337**Specification**

Recombinant Human NP-1 - Product Information**Recombinant Human NP-1 - Additional Information****Description**

Defensins (α and β) are cationic peptides with a broad spectrum of antimicrobial activity that comprise an important arm of the innate immune system. The α -defensins which include NP-1, NP-2 and NP-3, are distinguished from the β -defensins by the pairing of their three disulfide bonds. In addition to antimicrobial activity, NP-1 exhibits chemotactic activity on dendritic cells. NP-1 is expressed as the C-terminal portion of an inactive precursor protein, which also contains a 19 amino acid N-terminal signal sequence and a 45 amino acid polypeptide. NP-1 contains a six-cysteine motif that forms three intra-molecular disulfide bonds. Recombinant human NP-1 is a 3.4 kDa protein containing 30 amino acid residues.

Biological Activity

Determined by its ability to chemoattract immature dendritic cells using a concentration of 1.0-10.0 ng/ml.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is <0.1 ng/ μ g of protein (<1EU/ μ g).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C

Precautions

Recombinant Human NP-1 is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Human NP-1 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

Recombinant Human NP-1 - Images