

Recombinant Human sOX40 Ligand

Catalog # PBG10347

Specification

Recombinant Human sOX40 Ligand - Product Information

Recombinant Human sOX40 Ligand - Additional Information

Description

OX40L, a member of the TNF superfamily of structurally related proteins, exists primarily as a type II membrane bound, non-covalently linked homotrimeric protein. It is expressed on antigen presenting cells (APCs), such as dendritic cells and activated B-cells, and also on various other cells such as vascular endothelial cells, mast cells, and natural killer cells. OX40L signals specifically through the OX40 receptor, which is expressed predominantly on CD4+T cells but also on certain activated CD8+T cells. OX40/OX40L functions as a costimulatory signal, which is required for a productive interaction between antigen presenting cells and their target T-cells. It enhances cell proliferation and survival, and increases expression of RANTES, IL-2, IL-3, and IFNy. OX40/OX40L signaling plays an important role in immuno-regulatory communication, enabling the immune system to distinguish between "friend vs. foe" during activation; a mechanism typically termed immuno-tolerance. Recombinant OX40L is a glycosylated 133 amino acid protein corresponding to the extracellular TNF homologous domain of the full length transmembrane protein. It migrates with an apparent molecular mass of 15.5 – 25.0 kDa on SDS-PAGE.

BiologicalActivity

Determined by its ability to stimulate IL-8 production by human PBMC using a concentration range of 30-100 ng/ml. Note: Results may vary with PBMC donors.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is $<0.1 \text{ ng}/\mu\text{g}$ of protein ($<1\text{EU}/\mu\text{g}$).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C


Precautions

Recombinant Human sOX40 Ligand is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Human sOX40 Ligand - Protocols

Provided below are standard protocols that you may find useful for product applications.

Western Blot





• Blocking Peptides

- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Recombinant Human sOX40 Ligand - Images