

Recombinant Murine SDF-1 β (CXCL12)
Catalog # PBG10407**Specification**

Recombinant Murine SDF-1 β (CXCL12) - Product Information**Recombinant Murine SDF-1 β (CXCL12) - Additional Information****Description**

SDF-1 α and β are stromal derived CXC chemokines, and signal through the CXCR4 receptor. SDF-1 α and β chemoattract B and T cells, and have been shown to induce migration of CD34+ stem cells. Additionally, the SDF-1 proteins exert HIV suppressive activity in cells expressing the CXCR4 receptor. Recombinant murine SDF-1 β is a 8.5 kDa protein containing 72 amino acid residues.

Biological Activity

Determined by its ability to chemoattract human monocytes using a concentration range of 50.0-100.0 ng/ml.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is <0.1 ng/ μ g of protein (<1EU/ μ g).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C

Precautions

Recombinant Murine SDF-1 β (CXCL12) is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Murine SDF-1 β (CXCL12) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Recombinant Murine SDF-1 β (CXCL12) - Images

