

Recombinant Human TGF-β2 (mammalian derived)

Catalog # PBG10442

Specification

Recombinant Human TGF-β2 (mammalian derived) - Product Information

Recombinant Human TGF-β2 (mammalian derived) - Additional Information

Description

The three mammalian isoforms of TGF- β 1, TGF- β 1, β 2, β 3, signal through the same receptor and elicit similar biological responses. They are multifunctional cytokines that regulate cell proliferation, growth, differentiation and motility as well as synthesis and deposition of the extracellular matrix. They are involved in various physiological processes including embryogenesis, tissue remodeling and wound healing. They are secreted predominantly as latent complexes which are stored at the cell surface and in the extracellular matrix. The release of biologically active TGF- β 1 isoform from a latent complex involves proteolytic processing of the complex and /or induction of conformational changes by proteins such as thrombospondin-1. TGF- β 2 has been shown to exert suppressive effects on IL-2 dependent T-cell growth, and may also have an autocrine function in enhancing tumor growth by suppressing immuno-surveillance of tumor development. Recombinant human TGF- β 2 is a 25.0 kDa protein composed of two identical 112 amino acid polypeptide chains linked by a single disulfide bond.

BiologicalActivity

Determined by its ability to inhibit the M IL-4-dependent proliferation of M HT-2 cells. The ED₅₀ was found to be \le 0.2 ng/ml, corresponding to a specific activity of \ge 5 x 10⁶ units/mg.

Authenticity

Verified by N-terminal and Mass Spectrometry analyses (when applicable).

Endotoxin

Endotoxin level is $<0.1 \text{ ng}/\mu\text{g}$ of protein ($<1\text{EU}/\mu\text{g}$).

Protein Content

Verified by UV Spectroscopy and/or SDS-PAGE gel.

Storage

-20°C


Precautions

Recombinant Human TGF- β 2 (mammalian derived) is for research use only and not for use in diagnostic or therapeutic procedures.

Recombinant Human TGF-β2 (mammalian derived) - Protocols

Provided below are standard protocols that you may find useful for product applications.

Western Blot





• Blocking Peptides

- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Recombinant Human TGF- β 2 (mammalian derived) - Images