

Caspase-11 Blocking Peptide
Catalog # PBV10007b**Specification**

Caspase-11 Blocking Peptide - Product Information

Primary Accession	P70343
Other Accession	NM_007609
Gene ID	12363
Calculated MW	42742

Caspase-11 Blocking Peptide - Additional Information**Gene ID** 12363**Application & Usage**

The peptide is used for blocking the antibody activity of active Caspase-11. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30 minutes at 37°C

Other Names

Caspase-4, CASP-4, 3.4.22.64, Caspase-11, CASP-11, Protease ICH-3, Caspase-4 subunit p10, Caspase-4 subunit p20, Casp4, Casp11, Caspl, Ich3

Target/Specificity

Caspase-11

Formulation

50 µg (0.2 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 0.1% BSA and 0.02% thimerosal

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

Caspase-11 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

Caspase-11 Blocking Peptide - Protein Information**Name** Casp4 {ECO:0000312|MGI:MGI:107700}**Function**

Inflammatory caspase that acts as the effector of the non- canonical inflammasome by mediating lipopolysaccharide (LPS)-induced pyroptosis (PubMed:<a

[22002608](http://www.uniprot.org/citations/22002608), PubMed: [23348507](http://www.uniprot.org/citations/23348507), PubMed: [23887873](http://www.uniprot.org/citations/23887873), PubMed: [24031018](http://www.uniprot.org/citations/24031018), PubMed: [25119034](http://www.uniprot.org/citations/25119034), PubMed: [30135078](http://www.uniprot.org/citations/30135078), PubMed: [37001519](http://www.uniprot.org/citations/37001519)). Also indirectly activates the NLRP3 and NLRP6 inflammasomes (PubMed: [26320999](http://www.uniprot.org/citations/26320999), PubMed: [30392956](http://www.uniprot.org/citations/30392956), PubMed: [37001519](http://www.uniprot.org/citations/37001519)). Acts as a thiol protease that cleaves a tetrapeptide after an Asp residue at position P1: catalyzes cleavage of CGAS and GSDMD (PubMed: [26375003](http://www.uniprot.org/citations/26375003), PubMed: [28314590](http://www.uniprot.org/citations/28314590), PubMed: [30392956](http://www.uniprot.org/citations/30392956)). In contrast to its human ortholog, does not cleave IL18 (PubMed: [37993712](http://www.uniprot.org/citations/37993712), PubMed: [37993714](http://www.uniprot.org/citations/37993714)). Effector of the non-canonical inflammasome independently of NLRP3 inflammasome and CASP1: the non-canonical inflammasome promotes pyroptosis through GSDMD cleavage without involving secretion of cytokine IL1B and IL18 (PubMed: [22002608](http://www.uniprot.org/citations/22002608), PubMed: [22895188](http://www.uniprot.org/citations/22895188), PubMed: [23348507](http://www.uniprot.org/citations/23348507), PubMed: [23887873](http://www.uniprot.org/citations/23887873), PubMed: [24031018](http://www.uniprot.org/citations/24031018), PubMed: [26320999](http://www.uniprot.org/citations/26320999), PubMed: [26375003](http://www.uniprot.org/citations/26375003), PubMed: [30135078](http://www.uniprot.org/citations/30135078), PubMed: [30589883](http://www.uniprot.org/citations/30589883)). In the non-canonical inflammasome, CASP4/CASP11 is activated by direct binding to the lipid A moiety of LPS without the need of an upstream sensor (PubMed: [22002608](http://www.uniprot.org/citations/22002608), PubMed: [23348507](http://www.uniprot.org/citations/23348507), PubMed: [25119034](http://www.uniprot.org/citations/25119034), PubMed: [37001519](http://www.uniprot.org/citations/37001519)). LPS-binding promotes CASP4/CASP11 activation and CASP4/CASP11-mediated cleavage of GSDMD, followed by pyroptosis of infected cells and their extrusion into the gut lumen (PubMed: [22002608](http://www.uniprot.org/citations/22002608), PubMed: [23348507](http://www.uniprot.org/citations/23348507), PubMed: [25119034](http://www.uniprot.org/citations/25119034)). Also indirectly promotes secretion of mature cytokines (IL1A, IL18 and HMGB1) downstream of GSDMD-mediated pyroptosis via activation of the NLRP3 and NLRP6 inflammasomes (By similarity). Involved in NLRP3-dependent CASP1 activation and IL1B and IL18 secretion in response to non-canonical activators, such as UVB radiation or cholera enterotoxin (PubMed: [26320999](http://www.uniprot.org/citations/26320999)). Involved in NLRP6 inflammasome-dependent activation in response to lipoteichoic acid (LTA), a cell-wall component of Gram- positive bacteria, which leads to CASP1 activation and IL1B and IL18 secretion (PubMed: [30392956](http://www.uniprot.org/citations/30392956)). Involved in LPS-induced IL6 secretion; this activity may not require caspase enzymatic activity (By similarity). The non-canonical inflammasome is required for innate immunity to cytosolic, but not vacuolar, bacteria (PubMed: [23348507](http://www.uniprot.org/citations/23348507)). Plays a crucial role in the restriction of *S.typhimurium* replication in colonic epithelial cells during infection (PubMed: [25121752](http://www.uniprot.org/citations/25121752), PubMed: [26375003](http://www.uniprot.org/citations/26375003), PubMed: [34671164](http://www.uniprot.org/citations/34671164)). Pyroptosis limits bacterial replication, while cytokine secretion promotes the recruitment and

activation of immune cells and triggers mucosal inflammation (PubMed:25121752). May also act as an activator of adaptive immunity in dendritic cells, following activation by oxidized phospholipid 1- palmitoyl-2-arachidonoyl- sn-glycero-3-phosphorylcholine, an oxidized phospholipid (oxPAPC) (PubMed:27103670). Cleavage of GSDMD is not strictly dependent on the consensus cleavage site but depends on an exosite interface on CASP4/CASP11 that recognizes and binds the Gasdermin-D, C-terminal (GSDMD-CT) part (PubMed:32109412, PubMed:32554464). In contrast, it does not directly process IL1B (PubMed:8702803, PubMed:9038361). During non-canonical inflammasome activation, cuts CGAS and may play a role in the regulation of antiviral innate immune activation (PubMed:28314590).

Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:P49662}. Cytoplasm. Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P49662}; Peripheral membrane protein {ECO:0000250|UniProtKB:P49662}; Cytoplasmic side {ECO:0000250|UniProtKB:P49662}. Mitochondrion {ECO:0000250|UniProtKB:P49662}. Inflammasome. Secreted {ECO:0000250|UniProtKB:P49662} Note=Predominantly localizes to the endoplasmic reticulum (ER) Association with the ER membrane requires TMEM214. Released in the extracellular milieu by keratinocytes following UVB irradiation {ECO:0000250|UniProtKB:P49662}

Tissue Location

Widely expressed, including in thymus, lung and spleen (at protein level). Very low levels, if any, in the brain

Caspase-11 Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Caspase-11 Blocking Peptide - Images