

c-FLIP Blocking Peptide
Catalog # PBV10023b**Specification**

c-FLIP Blocking Peptide - Product Information

Primary Accession	O15519
Gene ID	8837
Calculated MW	55344

c-FLIP Blocking Peptide - Additional Information**Gene ID** 8837**Application & Usage**

The peptide is used for blocking the antibody activity of FLIP. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Other Names

CASP8 and FADD-like apoptosis regulator, Caspase homolog, CASH, Caspase-eight-related protein, Casper, Caspase-like apoptosis regulatory protein, CLARP, Cellular FLICE-like inhibitory protein, c-FLIP, FADD-like antiapoptotic molecule 1, FLAME-1, Inhibitor of FLICE, I-FLICE, MACH-related inducer of toxicity, MRIT, Usurpin, CASP8 and FADD-like apoptosis regulator subunit p43, CASP8 and FADD-like apoptosis regulator subunit p12, CFLAR, CASH, CASP8AP1, CLARP, MRIT

Target/Specificity

c-FLIP

Formulation

50 µg (0.2 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

c-FLIP Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

c-FLIP Blocking Peptide - Protein Information**Name** CFLAR**Synonyms** CASH, CASP8AP1, CLARP, MRIT

Function

Apoptosis regulator protein which may function as a crucial link between cell survival and cell death pathways in mammalian cells. Acts as an inhibitor of TNFRSF6 mediated apoptosis. A proteolytic fragment (p43) is likely retained in the death-inducing signaling complex (DISC) thereby blocking further recruitment and processing of caspase-8 at the complex. Full length and shorter isoforms have been shown either to induce apoptosis or to reduce TNFRSF-triggered apoptosis. Lacks enzymatic (caspase) activity.

Tissue Location

Widely expressed. Higher expression in skeletal muscle, pancreas, heart, kidney, placenta, and peripheral blood leukocytes. Also detected in diverse cell lines. Isoform 8 is predominantly expressed in testis and skeletal muscle

c-FLIP Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

c-FLIP Blocking Peptide - Images