

Perforin Blocking Peptide
Catalog # PBV10073b**Specification**

Perforin Blocking Peptide - Product Information

Primary Accession	Q5FVS5
Other Accession	NP_059026
Gene ID	50669

Perforin Blocking Peptide - Additional Information

Application & Usage	The peptide is used for blocking the antibody activity of Perforin. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.
---------------------	--

Target/Specificity
Perforin**Formulation**
50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.**Reconstitution & Storage**
-20 °C**Background Descriptions****Precautions**
Perforin Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.**Perforin Blocking Peptide - Protein Information****Perforin Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

Perforin Blocking Peptide - Images