

BMF Blocking Peptide
Catalog # PBV10132b**Specification**

BMF Blocking Peptide - Product Information

Primary Accession	Q3V5X8
Other Accession	NP_001030110.1
Gene ID	362100

BMF Blocking Peptide - Additional Information

Application & Usage	The peptide is used for blocking the antibody activity of BMF. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.
---------------------	--

Target/Specificity
BMF**Formulation**
50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.**Reconstitution & Storage**
-20 °C**Background Descriptions****Precautions**
BMF Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.**BMF Blocking Peptide - Protein Information****BMF Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

BMF Blocking Peptide - Images