

PUMA/bbc3 Blocking Peptide
Catalog # PBV10135b**Specification**

PUMA/bbc3 Blocking Peptide - Product Information

Primary Accession	Q96PG8
Gene ID	27113
Calculated MW	26498

PUMA/bbc3 Blocking Peptide - Additional Information**Gene ID** 27113**Application & Usage**

The peptide is used for blocking the antibody activity of PUMA. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Other Names

Bcl-2-binding component 3, JFY-1, p53 up-regulated modulator of apoptosis, BBC3, PUMA

Target/Specificity

PUMA/bbc3

Formulation

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

PUMA/bbc3 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

PUMA/bbc3 Blocking Peptide - Protein Information**Name** BBC3**Synonyms** PUMA**Function**

[Isoform 3]: Does not affect cell growth.

Cellular Location

Note=Contrary to isoforms 1 and 2, isoform 3 does not localize to the mitochondria

PUMA/bbc3 Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PUMA/bbc3 Blocking Peptide - Images