

Cytochrome c Blocking Peptide
Catalog # PBV10143b**Specification**

Cytochrome c Blocking Peptide - Product Information**Cytochrome c Blocking Peptide - Additional Information****Application & Usage**

The peptide is used for blocking the antibody activity of active r-Cytochrome c. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30 minutes at 37°C

Target/Specificity

Cytochrome c

Formulation

50 µg (0.2 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 0.1% BSA and 0.02% thimerosal.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

Cytochrome c Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

Cytochrome c Blocking Peptide - Protein Information**Cytochrome c Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cytochrome c Blocking Peptide - Images

