

ALK peptide
Catalog # PBV10146b**Specification**

ALK peptide - Product Information

Primary Accession	O9UM73
Gene ID	238
Calculated MW	176442

ALK peptide - Additional Information**Gene ID** 238**Application & Usage**

The peptide is used for blocking the antibody activity of ALK. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Other Names

ALK tyrosine kinase receptor, 2.7.10.1, Anaplastic lymphoma kinase, CD246, ALK

Target/Specificity

ALK

Formulation

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

ALK peptide is for research use only and not for use in diagnostic or therapeutic procedures.

ALK peptide - Protein Information**Name** ALK {ECO:0000303|PubMed:9174053, ECO:0000312|HGNC:HGNC:427}**Function**

Neuronal receptor tyrosine kinase that is essentially and transiently expressed in specific regions of the central and peripheral nervous systems and plays an important role in the genesis and differentiation of the nervous system (PubMed:11121404, PubMed:11387242, PubMed:11387242, PubMed:11387242).

[16317043](http://www.uniprot.org/citations/16317043), PubMed: [17274988](http://www.uniprot.org/citations/17274988), PubMed: [30061385](http://www.uniprot.org/citations/30061385), PubMed: [34646012](http://www.uniprot.org/citations/34646012), PubMed: [34819673](http://www.uniprot.org/citations/34819673)). Also acts as a key thinness protein involved in the resistance to weight gain: in hypothalamic neurons, controls energy expenditure acting as a negative regulator of white adipose tissue lipolysis and sympathetic tone to fine-tune energy homeostasis (By similarity). Following activation by ALKAL2 ligand at the cell surface, transduces an extracellular signal into an intracellular response (PubMed: [30061385](http://www.uniprot.org/citations/30061385), PubMed: [33411331](http://www.uniprot.org/citations/33411331), PubMed: [34646012](http://www.uniprot.org/citations/34646012), PubMed: [34819673](http://www.uniprot.org/citations/34819673)). In contrast, ALKAL1 is not a potent physiological ligand for ALK (PubMed: [34646012](http://www.uniprot.org/citations/34646012)). Ligand-binding to the extracellular domain induces tyrosine kinase activation, leading to activation of the mitogen-activated protein kinase (MAPK) pathway (PubMed: [34819673](http://www.uniprot.org/citations/34819673)). Phosphorylates almost exclusively at the first tyrosine of the Y-x-x-x-Y-Y motif (PubMed: [15226403](http://www.uniprot.org/citations/15226403), PubMed: [16878150](http://www.uniprot.org/citations/16878150)). Induces tyrosine phosphorylation of CBL, FRS2, IRS1 and SHC1, as well as of the MAP kinases MAPK1/ERK2 and MAPK3/ERK1 (PubMed: [15226403](http://www.uniprot.org/citations/15226403), PubMed: [16878150](http://www.uniprot.org/citations/16878150)). ALK activation may also be regulated by pleiotrophin (PTN) and midkine (MDK) (PubMed: [11278720](http://www.uniprot.org/citations/11278720), PubMed: [11809760](http://www.uniprot.org/citations/11809760), PubMed: [12107166](http://www.uniprot.org/citations/12107166), PubMed: [12122009](http://www.uniprot.org/citations/12122009)). PTN-binding induces MAPK pathway activation, which is important for the anti-apoptotic signaling of PTN and regulation of cell proliferation (PubMed: [11278720](http://www.uniprot.org/citations/11278720), PubMed: [11809760](http://www.uniprot.org/citations/11809760), PubMed: [12107166](http://www.uniprot.org/citations/12107166)). MDK-binding induces phosphorylation of the ALK target insulin receptor substrate (IRS1), activates mitogen-activated protein kinases (MAPKs) and PI3-kinase, resulting also in cell proliferation induction (PubMed: [12122009](http://www.uniprot.org/citations/12122009)). Drives NF-kappa-B activation, probably through IRS1 and the activation of the AKT serine/threonine kinase (PubMed: [15226403](http://www.uniprot.org/citations/15226403), PubMed: [16878150](http://www.uniprot.org/citations/16878150)). Recruitment of IRS1 to activated ALK and the activation of NF-kappa-B are essential for the autocrine growth and survival signaling of MDK (PubMed: [15226403](http://www.uniprot.org/citations/15226403), PubMed: [16878150](http://www.uniprot.org/citations/16878150)).

Cellular Location

Cell membrane; Single-pass type I membrane protein Note=Membrane attachment is essential for promotion of neuron-like differentiation and cell proliferation arrest through specific activation of the MAP kinase pathway.

Tissue Location

Expressed in brain and CNS. Also expressed in the small intestine and testis, but not in normal lymphoid cells

ALK peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ALK peptide - Images