

**Cox-2 Blocking Peptide**  
**Catalog # PBV10150b****Specification**

---

**Cox-2 Blocking Peptide - Product Information****Cox-2 Blocking Peptide - Additional Information****Application & Usage**

The peptide is used for blocking the antibody activity of active Cox-2. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30 minutes at 37°C

**Target/Specificity**  
Cox-2**Formulation**

50 µg (0.2 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 0.1% BSA and 0.02% thimerosal.

**Reconstitution & Storage**  
-20 °C**Background Descriptions****Precautions**

Cox-2 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

**Cox-2 Blocking Peptide - Protein Information****Cox-2 Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Cox-2 Blocking Peptide - Images**

