

recombinant protein Murine Interleukin-27/p28 (IL-30)

Interleukin-30, IL-30, IL-27/p28, p28, Interleukin-27, Interleukin-27/p28, IL-27, Interleukin-27 sub Catalog # PBV10166r

Specification

recombinant protein Murine Interleukin-27/p28 (IL-30) - Product info

Primary Accession <u>Q8K3I6</u>

Calculated MW 23.7 kDa KDa

recombinant protein Murine Interleukin-27/p28 (IL-30) - Additional Info

Gene ID 246779
Gene Symbol IL27

Other Names

Interleukin-30, IL-30, IL-27/p28, p28, Interleukin-27, Interleukin-27/p28, IL-27, Interleukin-27 subunit α , IL-27 subunit α , IL-27-A, II27, II27a, IL-27p28.

Gene Source Mouse Source E. coli

Assay&Purity SDS-PAGE; ≥97% Assay2&Purity2 HPLC; ≥97%

Recombinant Yes
Results 50 ng/ml

Target/Specificity

IL-27/p28

Application Notes

Reconstitute in H_2O to a concentration of 0.1-0.5 mg/ml. The solution can then be diluted into other aqueous buffers

Format

Lyophilized protein

Storage

-20°C; Lyophilized from 10 mM NaP, pH 7.5

recombinant protein Murine Interleukin-27/p28 (IL-30) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety





• Cell Culture

recombinant protein Murine Interleukin-27/p28 (IL-30) - Images

recombinant protein Murine Interleukin-27/p28 (IL-30) - Background

The p28 subunit of IL-27 (IL-27/p28), also known as Interleukin-30 (IL-30), is a member of the IL-12 family of cytokines. When combined with EBI3 (Epstein-Barr virus induced gene 3), p28 forms a heterodimer known as IL-27. p28 is thought to be a pro-inflammatory cytokine inducing immuno-modulatory effects, but the details of these effects are currently unclear and research is focused on delineating its specific biological functions. Recombinant mouse p28 is a non-glycosylated protein, containing 207 amino acids, with a molecular weight of 23.7 kDa.