

TRAF3 Blocking Peptide
Catalog # PBV10235b**Specification**

TRAF3 Blocking Peptide - Product Information**TRAF3 Blocking Peptide - Additional Information****Application & Usage**

The peptide is used for blocking the antibody activity of TRAF3. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Target/Specificity
TRAF3**Formulation**

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage
-20 °C**Background Descriptions****Precautions**

TRAF3 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

TRAF3 Blocking Peptide - Protein Information**TRAF3 Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRAF3 Blocking Peptide - Images

