

HSD17B4/MPF-2 Blocking Peptide

Catalog # PBV10267b

Specification

HSD17B4/MPF-2 Blocking Peptide - Product Information

Primary Accession	<u>P51659</u>
Gene ID	3295
Calculated MW	79686

HSD17B4/MPF-2 Blocking Peptide - Additional Information

Gene ID 3295

Application & Usage

The peptide is used for blocking the antibody activity of HSD17B4/MPF-2. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Other Names

Peroxisomal multifunctional enzyme type 2, MFE-2, 17-beta-hydroxysteroid dehydrogenase 4, 17-beta-HSD 4, D-bifunctional protein, DBP, Multifunctional protein 2, MPF-2, (3R)-hydroxyacyl-CoA dehydrogenase, 1.1.1.n12, Enoyl-CoA hydratase 2, 4.2.1.107, 4.2.1.119, 3-alpha, 7-alpha, 12-alpha-trihydroxy-5-beta-cholest-24-enoyl-CoA hydratase, HSD17B4, EDH17B4

Target/Specificity HSD17B4/MPF-2

Formulation

 $50~\mu g$ (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage -20 °C

Background Descriptions

Precautions HSD17B4/MPF-2 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

HSD17B4/MPF-2 Blocking Peptide - Protein Information

Name HSD17B4 (HGNC:5213)

Synonyms EDH17B4, SDR8C1



Function

Bifunctional enzyme acting on the peroxisomal fatty acid beta-oxidation pathway. Catalyzes two of the four reactions in fatty acid degradation: hydration of 2-enoyl-CoA (trans-2-enoyl-CoA) to produce (3R)-3-hydroxyacyl-CoA, and dehydrogenation of (3R)-3- hydroxyacyl-CoA to produce 3-ketoacyl-CoA (3-oxoacyl-CoA), which is further metabolized by SCPx. Can use straight-chain and branched-chain fatty acids, as well as bile acid intermediates as substrates.

Cellular Location Peroxisome.

Tissue Location Present in many tissues with highest concentrations in liver, heart, prostate and testis

HSD17B4/MPF-2 Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

HSD17B4/MPF-2 Blocking Peptide - Images