

BIF Blocking Peptide
Catalog # PBV10328b**Specification**

BIF Blocking Peptide - Product Information

Primary Accession	O9Y371
Gene ID	51100
Calculated MW	40796

BIF Blocking Peptide - Additional Information**Gene ID** 51100**Application & Usage**

The peptide is used for blocking the antibody activity of BIF. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30 minutes at 37°C.

Other Names

Endophilin-B1, Bax-interacting factor 1, Bif-1, SH3 domain-containing GRB2-like protein B1, SH3GLB1, KIAA0491

Target/Specificity

BIF

Formulation

50 µg (0.2 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 0.1% BSA and 0.02% sodium azide.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

BIF Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

BIF Blocking Peptide - Protein Information**Name** SH3GLB1**Synonyms** KIAA0491**Function**

May be required for normal outer mitochondrial membrane dynamics (PubMed:<a

[15452144](http://www.uniprot.org/citations/15452144)). Required for coatamer-mediated retrograde transport in certain cells (By similarity). May recruit other proteins to membranes with high curvature. May promote membrane fusion (PubMed:[11604418](http://www.uniprot.org/citations/11604418)). Involved in activation of caspase-dependent apoptosis by promoting BAX/BAK1 activation (PubMed:[16227588](http://www.uniprot.org/citations/16227588)). Isoform 1 acts proapoptotic in fibroblasts (By similarity). Involved in caspase-independent apoptosis during nutrition starvation and involved in the regulation of autophagy. Activates lipid kinase activity of PI3KC3 during autophagy probably by associating with the PI3K complex II (PI3KC3-C2) (PubMed:[17891140](http://www.uniprot.org/citations/17891140)). Associated with PI3KC3-C2 during autophagy may regulate the trafficking of ATG9A from the Golgi complex to the peripheral cytoplasm for the formation of autophagosomes by inducing Golgi membrane tubulation and fragmentation (PubMed:[21068542](http://www.uniprot.org/citations/21068542)). Involved in regulation of degradative endocytic trafficking and cytokinesis, probably in the context of PI3KC3-C2 (PubMed:[20643123](http://www.uniprot.org/citations/20643123)). Isoform 2 acts antiapoptotic in neuronal cells; involved in maintenance of mitochondrial morphology and promotes neuronal viability (By similarity).

Cellular Location

Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein. Mitochondrion outer membrane; Peripheral membrane protein. Cytoplasmic vesicle, autophagosome membrane. Midbody. Note=Association with the Golgi apparatus depends on the cell type (By similarity). Following starvation colocalizes with ATG5 and LC3 autophagy-related protein(s) on autophagosomal membranes (PubMed:17891140). {ECO:0000250, ECO:0000269|PubMed:17891140}

Tissue Location

Highly expressed in heart, skeletal muscle, kidney and placenta. Detected at lower levels in brain, colon, thymus, spleen, liver, small intestine, lung and peripheral blood leukocytes

BIF Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BIF Blocking Peptide - Images