

BLAME Blocking Peptide
Catalog # PBV10420b**Specification**

BLAME Blocking Peptide - Product Information

Primary Accession	D3ZS24
Other Accession	NP_001099443
Gene ID	289237

BLAME Blocking Peptide - Additional Information**Application & Usage**

The peptide is used for blocking the antibody activity of BLAME. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

Target/Specificity
BLAME**Formulation**

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

Reconstitution & Storage
-20 °C**Background Descriptions****Precautions**

BLAME Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

BLAME Blocking Peptide - Protein Information**BLAME Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

BLAME Blocking Peptide - Images