

**Swi6 Blocking Peptide**  
**Catalog # PBV10510b****Specification**

---

**Swi6 Blocking Peptide - Product Information**

|                   |                        |
|-------------------|------------------------|
| Primary Accession | <a href="#">O1ZZU3</a> |
| Gene ID           | <b>375757</b>          |
| Calculated MW     | <b>26739</b>           |

**Swi6 Blocking Peptide - Additional Information****Gene ID** 375757**Application & Usage**

The peptide is used for blocking the antibody activity of Swi6. It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30-60 minutes at 37°C.

**Other Names**

DNA repair protein SWI5 homolog, HBV DNAPTP1-transactivated protein A, Protein SAE3 homolog, SWI5, C9orf119, SAE3

**Target/Specificity**

Swi6

**Formulation**

50 µg (0.5 mg/ml) in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA and 0.02% thimerosal.

**Reconstitution & Storage**

-20 °C

**Background Descriptions****Precautions**

Swi6 Blocking Peptide is for research use only and not for use in diagnostic or therapeutic procedures.

**Swi6 Blocking Peptide - Protein Information****Name** SWI5**Synonyms** C9orf119, SAE3**Function**

Component of the SWI5-SFR1 complex, a complex required for double-strand break repair via

homologous recombination.

**Cellular Location**

Nucleus.

**Swi6 Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Swi6 Blocking Peptide - Images**