

K63-linked Di-Ubiquitin recombinant protein
K63-linked Di-Ubiquitin
Catalog # PBV10651r**Specification**

K63-linked Di-Ubiquitin recombinant protein - Product info

Concentration	2.5
Calculated MW	17.139 kDa (Band migrates faster on gels) KDa

K63-linked Di-Ubiquitin recombinant protein - Additional Info

Assay&Purity	Western Blot; ≥95%
Assay2&Purity2	N/A;
Format	
Liquid	

Storage

-80°C; 2.5 mg/ml in 20 mM Tris-HCl, pH 7.5, 0.15 M NaCl and 1 mM EDTA.

K63-linked Di-Ubiquitin recombinant protein - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

K63-linked Di-Ubiquitin recombinant protein - Images**K63-linked Di-Ubiquitin recombinant protein - Background**

Poly-ubiquitination of target proteins through K63 has recently become the focus of intense study. The topology of this linkage type is quite different from polyubiquitin linked through lysine 48. Modification of proteins by K63-linked polyubiquitination has been implicated in, among other cellular processes, the regulation of the DNA damage response, endosomal sorting, autophagy of misfolded/aggregated proteins, and neurodegeneration. These di-ubiquitin chains are generated from the enzymatic linkage of wild-type ubiquitin through lysine 63. The most distal ubiquitin contains an arginine substitution for a lysine at position 63, limiting chain length.