

SDF-1 β /CXCL2, rat recombinant protein
Stromal-Cell Derived Factor-1, CXCL12, PBSF
Catalog # PBV10830r**Specification**

SDF-1 β /CXCL2, rat recombinant protein - Product info

Primary Accession [O9OZD1](#)
Calculated MW **8.4 kDa** **KDa**

SDF-1 β /CXCL2, rat recombinant protein - Additional Info

Gene ID **24772**
Gene Symbol **SDF-1**
Other Names
Stromal-Cell Derived Factor-1, CXCL12, PBSF

Gene Source **RAT**
Source **E. Coli**
Assay&Purity **SDS-PAGE; \geq 95%**
Assay2&Purity2 **HPLC;**
Recombinant **Yes**
Sequence **KPVSLSYRCP CRFFESHVAR ANVKHLKILN
TPNCALQIVA RLKSNNRQVC IDPKLKWIQE
YLDKALNKRL KM**

Target/Specificity
CXCL2

Application Notes

Centrifuge the vial prior to opening. Reconstitute in water to a concentration of 0.1-1.0 mg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein (example 0.1% BSA) and store in working aliquots at -20°C to -80°C.

Format

Lyophilized powder

Storage

-20°C; Sterile filtered through a 0.2 micron filter. Lyophilized with no additives

SDF-1 β /CXCL2, rat recombinant protein - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SDF-1 β /CXCL2, rat recombinant protein - Images**SDF-1 β /CXCL2, rat recombinant protein - Background**

SDF-1 alpha and beta are stromal derived CXC chemokines, and signal through the CXCR4 receptor. SDF-1alpha and beta chemoattract B and T cells, and have been shown to induce migration of CD34+ stem cells. Additionally, the SDF-1 proteins exert HIV suppressive activity in cells expressing the CXCR4 receptor. Recombinant rat SDF-1 β is an 8.4 kDa protein containing 72 amino acid residues.