

Human CellExp CD84 / SLAMF5, human recombinant protein
CD84, SLAMF5, LY9B, SLAMF5, hCD84, mCD84
Catalog # PBV11001r**Specification**

Human CellExp CD84 / SLAMF5, human recombinant protein - Product infoPrimary Accession
Calculated MW[O9UIB8](#)

This protein is fused with 6×His tag at the C-terminus, has a calculated MW of 23.6 kDa. The predicted N-terminus is Lys 22. DTT-reduced Protein migrates as 35-45 kDa due to glycosylation. KDa

Human CellExp CD84 / SLAMF5, human recombinant protein - Additional InfoGene ID **8832**
Gene Symbol **CD84****Other Names**

CD84, SLAMF5, LY9B, SLAMF5, hCD84, mCD84

Gene Source **Human**
Source **HEK293 cells**
Assay&Purity **SDS-PAGE; ≥95%**
Assay2&Purity2 **N/A;**
Recombinant **Yes**
Results

Measured in a cell proliferation assay using PHA stimulated human T cells in the presence of anti-CD3. The ED50 for this effect is typically 2-6 µg/ml in the presence of anti-CD3 immobilized at 20 ng/ml.

Target/Specificity
CD84/SLAMF5**Application Notes**

Centrifuge the vial prior to opening. Reconstitute in sterile PBS, pH 7.4 to a concentration of 50 µg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 month. For extended storage, it is recommended to store at -20°C.

Format
Lyophilized**Storage**

-20°C; Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Generally 5-8% Mannitol or trehalose is added as a protectant before lyophilization.

Human CellExp CD84 / SLAMF5, human recombinant protein - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Human CellExp CD84 / SLAMF5, human recombinant protein - Images

Human CellExp CD84 / SLAMF5, human recombinant protein - Background

Leukocyte differentiation antigen CD84 is also known as SLAM family member 5 (SLAMF5), which belongs to immunoglobulin (Ig) superfamily. CD84 / SLAMF5 contain one Ig-like C2-type (immunoglobulin-like) domain. CD84 plays a role as adhesion receptor functioning by homophilic interactions and by clustering. CD84 / SLAMF5 increases proliferative responses of activated T-cells and SH2D1A/SAP does not seem to be required for this process. Homophilic interactions enhance interferon gamma/IFNG secretion in lymphocytes and induce platelet stimulation via a SH2D1A/SAP-dependent pathway.

Human CellExp CD84 / SLAMF5, human recombinant protein - References

de la Fuente M.A., et al. Blood 90:2398-2405(1997).
Krause S.W., et al. Biochem. J. 346:729-736(2000).
Palou E., et al. Tissue Antigens 55:118-127(2000).
Halleck A., et al. Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).