

Human CellExp CD62E/E-Selectin, human recombinant protein SELE, RP1-117P20.2, CD62E, ELAM1, ESEL, LECAM2, E-Selectin Catalog # PBV11055r

Specification

Human CellExp CD62E/E-Selectin, human recombinant protein - Product info

Primary Accession P16581

Calculated MW

This protein is fused with a 6×his tag at

C-terminus and has a calculated MW of 60 kDa expressed. The predicted N-terminus is Trp22. Protein migrates as 110 kDa in

reduced SDS-PAGE resulting from

glycosylation. KDa

Human CellExp CD62E/E-Selectin, human recombinant protein - Additional Info

Gene ID 6401
Gene Symbol SELE

Other Names

SELE, RP1-117P20.2, CD62E, ELAM1, ESEL, LECAM2, E-Selectin

Gene Source

Source

Assay&Purity

Human

HEK293 cells

SDS-PAGE; ≥95%

Assay2&Purity2
Recombinant

N/A;

Results Measured by the ability of the immobilized

protein to support the adhesion of U937 human histiocytic lymphoma cells. When 5 x 10^4 cells/well are added to human E-Selectin coated plates (2 μ g/mL with 100 μ L/well), >80% cells will adhere after 1

hour incubation at 37°C.

Target/Specificity CD62E/E-Selectin

Application Notes

Centrifuge the vial prior to opening. Reconstitute in sterile PBS, pH 7.4 to a concentration of 50 μ g/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 month. For extended storage, it is recommended to store at -20°C.

Format

Lyophilized

Storage

-20°C; Lyophilized from 0.22 μm filtered solution in PBS, pH7.4. Normally Mannitol or Trehalose is added as protectants before lyophilization.



Human CellExp CD62E/E-Selectin, human recombinant protein - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Human CellExp CD62E/E-Selectin, human recombinant protein - Images

Human CellExp CD62E/E-Selectin, human recombinant protein - Background

E-Selectin, also known as CD62 antigen-like family member E (CD62E), endothelial-leukocyte adhesion molecule 1 (ELAM-1), or leukocyte-endothelial cell adhesion molecule 2 (LECAM2), a member of the Selectin family, is a 107-115 kDa cell surface glycoprotein. It is transiently expressed on vascular endothelial cells in response to IL-1 β and TNF α . E selectin has a cassette structure: an N-terminal, C-type lectin domain, an EGF (epidermal-growth-factor)-like domain, 6 Sushi domain (SCR repeat) units, a transmembrane domain (TM) and an intracellular cytoplasmic tail (cyto). During inflammation, E-selectin plays an important part in recruiting leukocytes to the site of injury. The local release of cytokines IL-1 and TNF- α by damaged cells induces the over-expression of E-selectin on endothelial cells of nearby blood vessels. E-selectin mediates the adhesion of tumor cells to endothelial cells, by binding to E-selectin ligands expressed by neutrophils, monocytes, eosinophils, memory-effector T-like lymphocytes, natural killer cells or cancer cells. Furthermore, a number of studies have reported that levels of E-Selectin may be elevated in subjects with a variety of pathological conditions.

Human CellExp CD62E/E-Selectin, human recombinant protein - References

Hession C.,et al.Proc. Natl. Acad. Sci. U.S.A. 87:1673-1677(1990). Bevilacqua M.P.,et al.Science 243:1160-1165(1989). Collins T.,et al.J. Biol. Chem. 266:2466-2473(1991). Gregory S.G.,et al.Nature 441:315-321(2006). Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.