10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Human CellExp PD-1 /PDCD1, human recombinant protein

PD, PD 1, PD 1 protein, PD 1 recombinant protein, human PD 1 protein, human recombinant PD 1 protein

Catalog # PBV11119r

Specification

Human CellExp PD-1 /PDCD1, human recombinant protein - Product info

Primary Accession	Q15116
Calculated MW	**This protein is fused with a polyhistidine tag at the C-terminus, and has a calculated MW of 17.3 kDa. The predicted N-terminus is Pro 21. DTT-reduced Protein migrates as 38-42 kDa in SDS-PAGE due to glycosylation. KDa**

Human CellExp PD-1 /PDCD1, human recombinant protein - Additional Info

Gene ID	**5133**
Gene Symbol	**PDCD1**

Other Names
PDCD1, PD1, CD279, SLEB2, hPD-1, hPD-l

Gene Source	**Human**
Source	**HEK293 cells**
Assay&Purity	**SDS-PAGE; ≥95%**
Assay2&Purity2	**N/A;**
Recombinant	**Yes**
Results	**Measured by its binding ability in a functional ELISA. Immobilized rhPD1/PDCD1 at 2 μg/ml (100 μL/well) can bind rh PD-L1 / B7-H1 Fc Chimera with a linear range of 20-250 ng/mL.**

Target/Specificity
PD-1/PDCD1

Application Notes
Centrifuge the vial prior to opening. Reconstitute in sterile PBS, pH 7.4 to a concentration of 50 μg/ml. Do not vortex. This solution can be stored at 2-8°C for up to 1 month. For extended storage, it is recommended to store at -20°C.

Format
Lyophilized

Storage
-20°C; Lyophilized from 0.22 μm filtered solution in PBS, pH 7.4. Normally Mannitol or Trehalose is added as protectants before lyophilization.

Human CellExp PD-1 /PDCD1, human recombinant protein - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Human CellExp PD-1 /PDCD1, human recombinant protein - Images

Human CellExp PD-1 /PDCD1, human recombinant protein - Background

Programmed cell death protein 1 (PD-1) is also known as CD279 and PDCD1, is a type I membrane protein and is a member of the extended CD28/CTLA-4 family of T cell regulators. PDCD1 is expressed on the surface of activated T cells, B cells, macrophages, myeloid cells and a subset of thymocytes. PD-1 has two ligands, PD-L1 and PD-L2, which are members of the B7 family. PD-L1 is expressed on almost all murine tumor cell lines, including PA1 myeloma, P815 mastocytoma, and B16 melanoma upon treatment with IFN-γ. PD-L2 expression is more restricted and is expressed mainly by DCs and a few tumor lines. PD1 inhibits the T-cell proliferation and production of related cytokines including IL-1, IL-4, IL-10 and IFN-γ by suppressing the activation and transduction of PI3K/AKT pathway. In addition, coligation of PD1 inhibits BCR-mediating signal by dephosphorylating key signal transducer. In vitro, treatment of anti-CD3 stimulated T cells with PD-L1-Ig results in reduced T cell proliferation and IFN-γ secretion. Monoclonal antibodies targeting PD-1 that boost the immune system are being developed for the treatment of cancer. This protein is suitable for use in protein studies such as protein structure analysis and protein-protein interactions. It can also be used as an immunogen, as a protein standard, or in cell biology research applications.

Human CellExp PD-1 /PDCD1, human recombinant protein - References

Shinohara T.,et al.Genomics 23:704-706(1994).
Finger L.R.,et al.Gene 197:177-187(1997).
Finger L.R.,et al.Gene 203:253-253(1997).
Prokunina L.,et al.Nat. Genet. 32:666-669(2002).
He X.,et al.Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.