

PKC delta, Active recombinant protein
PKC, Protein kinase C delta
Catalog # PBV11314r**Specification**

PKC delta, Active recombinant protein - Product info

Primary Accession	Q05655
Concentration	0.1
Calculated MW	104.0 kDa KDa

PKC delta, Active recombinant protein - Additional Info

Gene ID	5580
Gene Symbol	PRKCD
Other Names	
PKC, Protein kinase C delta	

Source	Baculovirus (Sf9 insect cells)
Assay&Purity	SDS-PAGE; ≥80%
Assay2&Purity2	HPLC;
Recombinant	Yes
Format	
Liquid	

Storage

-80°C; Recombinant protein in storage buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.25 mM DTT, 0.1 mM EGTA, 0.1 mM EDTA, 0.1 mM PMSF, 25% glycerol).

PKC delta, Active recombinant protein - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PKC delta, Active recombinant protein - Images**PKC delta, Active recombinant protein - Background**

Protein kinase C delta (PKC delta) is a member of the protein kinase C (PKC) family of serine-threonine kinases. It is a 79 kd protein kinase that shows strict dependence on the presence of phospholipids, but shows no activation by Ca²⁺ (1). Phosphatidylinositol is the most potent activator of PKC delta. Apparent kinetic constants for synthetic oligopeptides (MBP4-14, EGFR

peptide and epsilon-peptide) are quite different between PKC delta and other PKCs. Northern blot analysis indicated that PKC delta is widely distributed in almost all the tissues and is a major isoform of PKC expressed in hemopoietic cells (2). PKC delta is involved in fundamental cellular functions regulated by diacylglycerols and mimicked by phorbol esters. PKC delta is partially associated with the insoluble fraction in cells even in the absence of phorbol 12-myristate 13-acetate (PMA). Upon PMA stimulation, both it translocate to the insoluble fraction of cell homogenates (3). Overexpression of PKC-delta induces significant changes in morphology and causes the cells to grow more slowly and to a decreased cell density in confluent cultures. These changes are accentuated by treatment with PMA. None of the PKC-delta overexpressers grow in soft agar with or without PMA.