

GRO- α /KC/CINC-1/CXCL1
Catalog # PVGS1099**Specification**

GRO- α /KC/CINC-1/CXCL1 - Product Information

Primary Accession [P14095](#)
Species
Rat

Sequence
Ala25-Lys96

Purity
> 97% as analyzed by SDS-PAGE
> 97% as analyzed by HPLC

Endotoxin Level
< 1 EU/ μ g of protein by LAL method

Biological Activity
Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using rat neutrophils is in a concentration range of 10.0-100.0 ng/ml.

Expression System
E. coli

Theoretical Molecular Weight
7.8 kDa

Formulation **Lyophilized from a 0.2 μ m filtered solution in 20 mM PB, pH 7.4, 150 mM NaCl.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

GRO- α /KC/CINC-1/CXCL1 - Additional Information

Gene ID 81503

Other Names
Growth-regulated alpha protein, C-X-C motif chemokine 1, Cytokine-induced neutrophil chemoattractant 1, CINC-1, Platelet-derived growth factor-inducible protein KC, Cxcl1, Cinc1, Gro, Scyb1

Target Background

GRO- α /KC/CINC-1/CXCL1 has chemotactic activity for neutrophils. It may play a role in inflammation and exerts its effects on endothelial cells in an autocrine fashion. All three isoforms of GRO are CXC chemokines that can signal through the CXCR1 or CXCR2 receptors. GRO expression is inducible by serum or PDGF and/or by a variety of inflammatory mediators, such as IL-1 and TNF, in monocytes, fibroblasts, melanocytes and epithelial cells. In certain tumor cell lines, GRO is expressed constitutively.

GRO- α /KC/CINC-1/CXCL1 - Protein Information

Name Cxcl1

Synonyms Cinc1, Gro, Scyb1

Function

Has chemotactic activity for neutrophils. Contributes to neutrophil activation during inflammation.

Cellular Location

Secreted.

Tissue Location

At least expressed in the lung and trachea.

GRO- α /KC/CINC-1/CXCL1 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GRO- α /KC/CINC-1/CXCL1 - Images