

**GRO-γ/CXCL3**  
**Catalog # PVGS1111****Specification**

---

**GRO-γ/CXCL3 - Product Information**

Primary Accession [P19876](#)  
**Species**  
Human

**Sequence**  
Ala35-Asn107

**Purity**  
> 97% as analyzed by SDS-PAGE<br>> 97% as analyzed by HPLC

**Endotoxin Level**  
< 1 EU/ μg of protein by LAL method

**Biological Activity**  
Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human CXCR2 transfected human 293 cells is in a concentration range of 10.0-100.0 ng/ml.

**Expression System**  
E. coli

**Theoretical Molecular Weight**  
7.9 kDa

Formulation **Lyophilized from a 0.2 μm filtered solution in 20 mM PB, pH 7.4, 50 mM NaCl.**

**Reconstitution**  
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

**Storage & Stability**  
Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

**GRO-γ/CXCL3 - Additional Information**

**Gene ID** 2921

**Other Names**  
C-X-C motif chemokine 3, GRO-gamma(1-73), Growth-regulated protein gamma, GRO-gamma, Macrophage inflammatory protein 2-beta, MIP2-beta, GRO-gamma(5-73), CXCL3, GRO3, GROG, SCYB3

**Target Background**

Chemokine (C-X-C motif) ligand 3 (CXCL3) is a small cytokine belonging to the CXC chemokine family that is also known as GRO3 oncogene (GRO3), GRO protein gamma (GROγ) and macrophage inflammatory protein-2-beta (MIP2b). CXCL3 controls migration and adhesion of monocytes and mediates its effect on its target cell by interacting with cell surface chemokine receptor CXCR2. It has been shown that CXCL3 regulates the migration of precursors of cerebellar granule neurons toward the internal layers of the cerebellum, during morphogenesis.

**GRO-γ/CXCL3 - Protein Information**

**Name** CXCL3

**Synonyms** GRO3, GROγ, SCYB3

**Function**

Ligand for CXCR2 (By similarity). Has chemotactic activity for neutrophils. May play a role in inflammation and exert its effects on endothelial cells in an autocrine fashion. In vitro, the processed form GRO-γ(5-73) shows a fivefold higher chemotactic activity for neutrophilic granulocytes.

**Cellular Location**

Secreted.

**GRO-γ/CXCL3 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**GRO-γ/CXCL3 - Images**