

Eotaxin/CCL11
Catalog # PVGS1121**Specification**

Eotaxin/CCL11 - Product Information

Primary Accession [P51671](#)
Species
Human

Sequence
Gly24-Pro97

Purity
> 97% as analyzed by SDS-PAGE
> 97% as analyzed by HPLC

Endotoxin Level
< 1 EU/ µg of protein by LAL method

Biological Activity
Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood eosinophils is in a concentration range of 0.1-10.0 ng/ml.

Expression System
E. coli

Theoretical Molecular Weight
8.4 kDa

Formulation **Lyophilized from a 0.2 µm filtered solution in 20 mM PB, pH 7.4, 150 mM NaCl.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

Eotaxin/CCL11 - Additional Information

Gene ID 6356

Other Names
Eotaxin, C-C motif chemokine 11, Eosinophil chemotactic protein, Small-inducible cytokine A11, CCL11, SCYA11

Target Background

CCL11 is a potent eosinophil chemoattractant that was originally purified from bronchoalveolar lavage fluid of guinea pigs sensitized by aerosol challenge with ovalbumin. Human CCL11 cDNA encodes a 97 amino acid residue precursor protein from which the aminoterminal 23 amino acid residues are cleaved to generate the 74 amino acid residue mature human CCL11. At the protein sequence level, mature human CCL11 is approximately 60% identical to mature mouse and guinea pig CCL11. Human CCL11 is chemotactic for eosinophils, but not mononuclear cells or neutrophils. The CC chemokine receptor 3 (CCR3) has now been identified to be a specific human CCL11 receptor. CCR3 has also been shown to serve as a cofactor for a restricted subset of primary HIV viruses and binding of CCL11 to CCR3 inhibited infection by the HIV isolates.

Eotaxin/CCL11 - Protein Information

Name CCL11

Synonyms SCYA11

Function

In response to the presence of allergens, this protein directly promotes the accumulation of eosinophils, a prominent feature of allergic inflammatory reactions (PubMed:8597956). Binds to CCR3 (PubMed:8631813).

Cellular Location

Secreted.

Eotaxin/CCL11 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Eotaxin/CCL11 - Images