

IFN- ω
Catalog # PVGS1165

Specification

IFN- ω - Product Information

Primary Accession [P05000](#)
Species
Human

Sequence
Cys24-Ser195

Purity
> 97% as analyzed by SDS-PAGE
> 97% as analyzed by HPLC

Endotoxin Level
< 1 EU/ μ g of protein by LAL method

Biological Activity
Fully biologically active when compared to standard. The ED₅₀ as determined by a chemotaxis bioassay using human TF-1 cells is less than 0.01 ng/ml, corresponding to a specific activity of > 1.0×10^8 IU/mg.

Expression System
E. coli

Theoretical Molecular Weight
20 kDa

Formulation **Lyophilized from a 0.2 μ m filtered solution in PBS, pH 7.4.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at -70°C or -20°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

IFN- ω - Additional Information

Gene ID 3467

Other Names
Interferon omega-1, Interferon alpha-II-1, IFNW1

Target Background

Interferon-Omega (IFN- ω) coded by IFNW1 gene in human, is a member of the type I interferon family, which includes IFN- α , IFN- β , and IFN- ω . The IFNAR-1/IFNAR-2 receptor complex can help with the signal transduction, followed by the antiviral or the antiproliferative actions. IFN- ω is derived from IFN- α/β and shares 75% sequence with IFN- α . It has two intramolecular disulfide bonds which are crucial for its activities. Mire-Sluis et al have described bioassays for IFN- α , IFN- β , and IFN- ω that exploit the ability of these factors to inhibit proliferation of TF-1 cells induced by GM-CSF. The bioassays can be used also with Epo and TF-1 cells, or Epo and Epo-transfected UT-7 cells.

IFN- ω - Protein Information

Name IFNW1

Cellular Location

Secreted.

IFN- ω - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IFN- ω - Images