

**G-CSF**  
**Catalog # PVGS1194****Specification**

---

**G-CSF - Product Information**

Primary Accession [Q8N4W3](#)  
**Species**  
Human

**Sequence**  
Thr27-Pro200

**Purity**  
> 95% as analyzed by SDS-PAGE  
> 95% as analyzed by HPLC

**Endotoxin Level**  
< 0.2 EU/ µg of protein by gel clotting method

**Biological Activity**  
ED<sub>50</sub> < 0.1 ng/ml, determined by the dose-dependent stimulation of the proliferation of murine M-NFS-60 cells, corresponding to a specific activity of > 1.0 × 10<sup>7</sup> units/mg.

**Expression System**  
CHO

Formulation **Lyophilized after extensive dialysis against PBS.**

**Reconstitution**  
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH<sub>2</sub>O or PBS up to 100 µg/ml.

**Storage & Stability**  
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

**G-CSF - Additional Information**

**Target Background**  
Human Granulocyte Colony Stimulating Factor (G-CSF) contains internal disulfide bonds. Among the family of colony-stimulating factors, Granulocyte Colony Stimulating Factor (G-CSF) is the most potent inducer of terminal differentiation to granulocytes and macrophages of leukemic myeloid cell lines. The synthesis of Granulocyte Colony Stimulating Factor (G-CSF) can be induced by bacterial endotoxins, TNF, Interleukin-1 and GM-CSF. Prostaglandin E2 inhibits the synthesis of Granulocyte Colony Stimulating Factor (G-CSF). In epithelial, endothelial, and fibroblastic cells, the secretion of Granulocyte Colony Stimulating Factor (G-CSF) is induced by Interleukin-17.

## **G-CSF - Protein Information**

## **G-CSF - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **G-CSF - Images**