

GRO- α /MGSA/CXCL1
Catalog # PVGS1247**Specification**

GRO- α /MGSA/CXCL1 - Product Information

Primary Accession [P09341](#)
Species
Human

Sequence
Ala35-Asn107

Purity
> 95% as analyzed by SDS-PAGE

Endotoxin Level
< 0.2 EU/ μ g of protein by gel clotting method

Biological Activity
Active, measured in a functional assay using HUVEC cells.

Expression System
CHO

Formulation **Lyophilized after extensive dialysis against PBS.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or PBS up to 100 μ g/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

GRO- α /MGSA/CXCL1 - Additional Information

Gene ID 2919

Other Names
Growth-regulated alpha protein, C-X-C motif chemokine 1, GRO-alpha(1-73), Melanoma growth stimulatory activity, MGSA, Neutrophil-activating protein 3, NAP-3, GRO-alpha(4-73), GRO-alpha(5-73), GRO-alpha(6-73), CXCL1, GRO, GRO1, GROA, MGSA, SCYB1

Target Background
GRO- α /MGSA/CXCL1 has chemotactic activity for neutrophils. It may play a role in inflammation and exerts its effects on endothelial cells in an autocrine fashion. All three isoforms of GRO are CXC chemokines that can signal through the CXCR1 or CXCR2 receptors. GRO expression is

inducible by serum or PDGF and/or by a variety of inflammatory mediators, such as IL-1 and TNF, in monocytes, fibroblasts, melanocytes and epithelial cells. In certain tumor cell lines, GRO is expressed constitutively.

GRO- α /MGSA/CXCL1 - Protein Information

Name CXCL1

Synonyms GRO, GRO1, GROA, MGSA, SCYB1

Function

Has chemotactic activity for neutrophils. May play a role in inflammation and exerts its effects on endothelial cells in an autocrine fashion. In vitro, the processed forms GRO-alpha(4-73), GRO-alpha(5-73) and GRO-alpha(6-73) show a 30-fold higher chemotactic activity.

Cellular Location

Secreted.

GRO- α /MGSA/CXCL1 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GRO- α /MGSA/CXCL1 - Images