

Epigen

Catalog # PVGS1320

Specification

Epigen - Product Information

Primary Accession
Species
Human

Q6UW88-2

Sequence

Ala24-Ala95, expressed with an N-terminal Met

Purity

> 95% as analyzed by SDS-PAGE

Endotoxin Level

< 0.2 EU/ μg of protein by gel clotting method

Biological Activity

ED₅₀ < 1.0 µg/ml, measured by a cell proliferation assay using 3T3 cells, corresponding to a specific activity of > 1.0 × 10³ units/mg.

Expression System

E. coli

Formulation

Lyophilized after extensive dialysis against PBS.

Reconstitution

It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH_2O up to $100 \mu g/ml$.

Storage & Stability

Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Epigen - Additional Information

Target Background

Epigen is a cytokine belonging to the Epidermal Growth Factor (EGF) superfamily, which also includes Epiregulin, Amphiregulin, Neuregulin 2- β , and Transforming Growth Factor α . The precursor of Epigen produced in tissues has 154 amino acids, and shares the characteristics of other members of EGF superfamily, including 3 disulfide bonds formed by 6 cysteines. Epigen is present in testis, heart, and liver, and it binds to EGF receptors with a much lower binding affinity than EGF. However, Epigen is more mitogenic than EGF. Epigen achieves its strong mitogenic potency by suppressing ligand-induced receptor inactivation. Unlike EGF, Epigen can also bind to EGF receptors in low pH conditions, helping its recycling. Therefore Epigen has anomalous potency due to its prolonged presence.





Tel: 858.875.1900 Fax: 858.875.1999

Epigen - Protein Information

Epigen - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Epigen - Images