

Leptin

Catalog # PVGS1329

Specification

Leptin - Product Information

Primary Accession **Species**Rat

P50596

Sequence

Val22-Cys167, expressed with an N-terminal Met

Purity

> 95% as analyzed by SDS-PAGE
br>> 95% as analyzed by HPLC

Endotoxin Level

< 0.2 EU/ μg of protein by gel clotting method

Biological Activity

ED₅₀ < 10.0 μ g/ml, measured by a cell proliferation assay using LoVo cells, corresponding to a specific activity of > 100.0 units/mg.

Expression System

E. coli

Formulation

Lyophilized after extensive dialysis against 50 mM Tris, pH 8.0.

Reconstitution

It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH_2O up to $100 \mu g/ml$.

Storage & Stability

Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

Leptin - Additional Information

Gene ID 25608

Other Names

Leptin, Obesity factor, Lep, Ob

Target Background

Leptin is a cytokine belonging to the Interleukin 6 family, and has a four-helix bundle structure. Leptin is encoded by the ob gene, and produced and secreted by white adipose tissue. The receptors of Leptin are Type I cytokine receptors, which exist in two different forms: a short form expressed in multiple tissues, and a long form expressed exclusively in the central nervous system



(CNS). Upon binding to Leptin, the receptors activate the JAK/STAT3 pathway and PI3K, and stimulate transcriptional programs that regulate feeding behavior, metabolic rate, endocrine axes, and glucose fluxes. The deficiency of Leptin in human and mouse causes morbid obesity, diabetes, and neuroendocrine anomalies. Leptin also has effects on reproduction and immunity. In summary, Leptin is a pivotal cytokine controlling energy balance, and as such has profound effects on human health.

Leptin - Protein Information

Name Lep


Synonyms Ob

Function

Key player in the regulation of energy balance and body weight control. Once released into the circulation, has central and peripheral effects by binding LEPR, found in many tissues, which results in the activation of several major signaling pathways (By similarity) (PubMed:11677594). In the hypothalamus, acts as an appetite-regulating factor that induces a decrease in food intake and an increase in energy consumption by inducing anorexinogenic factors and suppressing orexigenic neuropeptides, also regulates bone mass and secretion of hypothalamo-pituitary-adrenal hormones. In the periphery, increases basal metabolism, influences reproductive function, regulates pancreatic beta-cell function and insulin secretion, is pro-angiogenic for endothelial cell and affects innate and adaptive immunity (By similarity) (PubMed: 10482243, PubMed:11460888, PubMed:11677594, PubMed:9003024). In the arcuate nucleus of the hypothalamus, activates by depolarization POMC neurons inducing FOS and SOCS3 expression to release anorexigenic peptides and inhibits by hyperpolarization NPY neurons inducing SOCS3 with a consequent reduction on release of orexigenic peptides (By similarity) (PubMed:10482243, PubMed:9003024). In addition to its known satiety inducing effect, has a modulatory role in nutrient absorption. In the intestine, reduces glucose absorption by enterocytes by activating PKC and leading to a sequential activation of p38, PI3K and ERK signaling pathways which exerts an inhibitory effect on glucose absorption (By similarity). Acts as a growth factor on certain tissues, through the activation of different signaling pathways increases expression of genes involved in cell cycle regulation such as CCND1, via IAK2-STAT3 pathway, or VEGFA, via MAPK1/3 and PI3K-AKT1 pathways (By similarity). May also play an apoptotic role via JAK2- STAT3 pathway and up-regulation of BIRC5 expression (By similarity). Pro-angiogenic, has mitogenic activity on vascular endothelial cells and plays a role in matrix remodeling by regulating the expression of matrix metalloproteinases (MMPs) and tissue inhibitors of metalloproteinases (TIMPs) (PubMed:11460888). In innate immunity, modulates the activity and function of neutrophils by increasing chemotaxis and the secretion of oxygen radicals. Increases phagocytosis by macrophages and enhances secretion of pro-inflammatory mediators. Increases cytotoxic ability of NK cells (Probable). Plays a proinflammatory role, in synergy with IL1B, by inducing NOS2 which promotes the production of IL6, IL8 and Prostaglandin E2, through a signaling pathway that involves JAK2, PI3K, MAP2K1/MEK1 and MAPK14/p38 (By similarity). In adaptive immunity, promotes the switch of memory T- cells towards T helper-1 cell immune responses (By similarity). Increases CD4(+)CD25(-) T-cell proliferation and reduces autophagy during TCR (T-cell receptor) stimulation, through MTOR signaling pathway activation and BCL2 up-regulation (By similarity).

Cellular Location





Secreted.

Leptin - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Leptin - Images