

G-CSF
Catalog # PVGS1345**Specification**

G-CSF - Product Information

Primary Accession [P09920](#)
Species
Mouse

Sequence
Val31-Ala208

Purity
> 95% as analyzed by SDS-PAGE
> 95% as analyzed by HPLC

Endotoxin Level
< 0.2 EU/ µg of protein by gel clotting method

Biological Activity
ED₅₀ < 0.02 ng/ml, measured in a cell proliferation assay using NFS-60 cells.

Expression System
CHO

Formulation **Lyophilized after extensive dialysis against PBS.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or PBS up to 100 µg/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

G-CSF - Additional Information

Gene ID 12985

Other Names
Granulocyte colony-stimulating factor, G-CSF, Csf3, CsfG

Target Background
Granulocyte Colony-Stimulating Factor (G-CSF), also known as CSF-3 and MGI-1G, is a cytokine and hormone belonging to the IL-6 superfamily. It is expressed by monocytes, macrophages, endothelial cells, fibroblasts and bone marrow stroma. G-CSF stimulates the bone marrow to produce granulocytes and stem cells, and specifically stimulates the proliferation and differentiation of the neutrophilic granulocyte lineage. G-CSF has been used to stimulate white

blood cell production after chemotherapy. It has also been used to boost the number of hematopoietic stem cells after bone marrow transplantation.

G-CSF - Protein Information

Name Csf3

Synonyms Csf3

Function

Granulocyte/macrophage colony-stimulating factors are cytokines that act in hematopoiesis by controlling the production, differentiation, and function of 2 related white cell populations of the blood, the granulocytes and the monocytes-macrophages. This CSF induces granulocytes.

Cellular Location

Secreted.

G-CSF - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

G-CSF - Images