

**Adiponectin/Acrp30**  
**Catalog # PVGS1359****Specification**

---

**Adiponectin/Acrp30 - Product Information**Primary Accession [Q15848](#)**Species**  
Human**Sequence**  
Glu19-Asn244**Purity**  
> 95% as analyzed by SDS-PAGE  
> 95% as analyzed by HPLC**Endotoxin Level**  
< 0.2 EU/ µg of protein by gel clotting method**Biological Activity**  
ED<sub>50</sub> < 20.0 µg/ml, measured in a cell growth inhibition assay using M1 cells.**Expression System**  
CHO**Formulation** **Lyophilized after extensive dialysis against PBS.****Reconstitution**  
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH<sub>2</sub>O or PBS up to 100 µg/ml.**Storage & Stability**  
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.**Adiponectin/Acrp30 - Additional Information****Gene ID** 9370**Other Names**  
Adiponectin, 30 kDa adipocyte complement-related protein, Adipocyte complement-related 30 kDa protein, ACRP30, Adipocyte, C1q and collagen domain-containing protein, Adipose most abundant gene transcript 1 protein, apM-1, Gelatin-binding protein, ADIPOQ**Target Background**  
Adiponectin is an important adipokine involved in the control of fat metabolism and insulin sensitivity. It is synthesized exclusively by adipocytes and secreted into plasma. It antagonizes THF-α by negatively regulating its expression. It also inhibits endothelial NF-κB signaling

through a cAMP-dependent pathway. Adiponectin can form low molecular weight complexes (LMW), middle molecular weight complexes (MMW) and higher molecular weight complexes (HMW). These bind to various growth factors, such as HBEGF, PDGFB and FGF2, and play a role in cell growth, angiogenesis and tissue remodeling.

## **Adiponectin/Acrp30 - Protein Information**

**Name** ADIPOQ

### **Function**

Important adipokine involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Stimulates AMPK phosphorylation and activation in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Antagonizes TNF-alpha by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. Inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. May play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex, LMW, MMW or HMW.

### **Cellular Location**

Secreted.

### **Tissue Location**

Synthesized exclusively by adipocytes and secreted into plasma.

## **Adiponectin/Acrp30 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Adiponectin/Acrp30 - Images**