

VEGF-D
Catalog # PVGS1366**Specification**

VEGF-D - Product Information

Primary Accession [O43915](#)
Species
Human

Sequence
Phe89-Arg205

Purity
> 95% as analyzed by SDS-PAGE
> 95% as analyzed by HPLC

Endotoxin Level
< 0.2 EU/ µg of protein by gel clotting method

Biological Activity
ED₅₀ < 1.0 µg/ml, measured in a cell proliferation assay using HUVEC cells.

Expression System
CHO

Formulation **Lyophilized after extensive dialysis against PBS.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or PBS up to 100 µg/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

VEGF-D - Additional Information

Gene ID 2277

Other Names
Vascular endothelial growth factor D {ECO:0000312|HGNC:HGNC:3708}, VEGF-D, c-Fos-induced growth factor, FIGF, VEGFD (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=3708), FIGF

Target Background
Vascular Endothelial Growth Factor (VEGF)-D, also known as c-Fos-induced growth factor (FIGF), is a member of the PDGF/VEGF growth factor family. It is expressed highly in lung, heart and small

intestine, and at lower levels in skeletal muscle, colon and pancreas. It binds to VEGFR-2 and VEGFR-3 receptors and activates downstream signals. VEGF-D is a growth factor active in angiogenesis, lymphangiogenesis and endothelial cell growth. It is involved in many developmental and physiological processes including the formation of venous and lymphatic vascular systems during embryogenesis and the maintenance of differentiated lymphatic endothelium in adults. In tumor pathology, it has been reported to play a role in restructuring of lymphatic channels and regional lymph node metastasis.

VEGF-D - Protein Information

Name VEGFD ([HGNC:3708](#))

Synonyms FIGF

Function

Growth factor active in angiogenesis, lymphangiogenesis and endothelial cell growth, stimulating their proliferation and migration and also has effects on the permeability of blood vessels. May function in the formation of the venous and lymphatic vascular systems during embryogenesis, and also in the maintenance of differentiated lymphatic endothelium in adults. Binds and activates VEGFR-2 (KDR/FLK1) and VEGFR-3 (FLT4) receptors.

Cellular Location

Secreted.

Tissue Location

Highly expressed in lung, heart, small intestine and fetal lung, and at lower levels in skeletal muscle, colon, and pancreas

VEGF-D - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VEGF-D - Images