

**MPIF-1/CCL23**  
**Catalog # PVGS1426****Specification**

---

**MPIF-1/CCL23 - Product Information**

Primary Accession [P55773](#)  
**Species**  
Human

**Sequence**  
Arg22-Asn120

**Purity**  
> 98% as analyzed by SDS-PAGE

**Endotoxin Level**  
< 0.2 EU/ µg of protein by gel clotting method

**Biological Activity**  
The EC<sub>50</sub> value of human MPIF-1/CCL23 on Ca<sup>2+</sup> mobilization assay in CHO-K1/Ga15/hCCR1 cells (human Ga15 and human CCR1 stably expressed in CHO-K1 cells) is less than 2.0 µg/ml.

**Expression System**  
CHO

Formulation **Lyophilized after extensive dialysis against PBS.**

**Reconstitution**  
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH<sub>2</sub>O or PBS up to 100 µg/ml.

**Storage & Stability**  
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

**MPIF-1/CCL23 - Additional Information**

**Gene ID** 6368

**Other Names**  
C-C motif chemokine 23, CK-beta-8, CKB-8, Macrophage inflammatory protein 3, MIP-3, Myeloid progenitor inhibitory factor 1, MPIF-1, Small-inducible cytokine A23, CCL23(19-99), CCL23(22-99), CCL23(27-99), CCL23(30-99), CCL23, MIP3, MPIF1, SCYA23

**Target Background**  
Myeloid progenitor inhibitory factor 1 (MPIF-1), also known as Chemokine (C-C motif) ligand 23

(CCL23) is a small cytokine belonging to the CC chemokine family. MPIF-1 is predominantly expressed in lung and liver tissue, but is also found in bone marrow and placenta. It is also expressed in some cell lines of myeloid origin. It is highly chemotactic for resting T cells and monocytes and slightly chemotactic for neutrophils. MPIF-1 has been shown to inhibit colony formation of bone marrow myeloid immature progenitors. It has also been attributed to an inhibitory activity on hematopoietic progenitor cells. MPIF-1 is a ligand for the chemokine receptor CCR1.

## **MPIF-1/CCL23 - Protein Information**

**Name** CCL23

**Synonyms** MIP3, MPIF1, SCYA23

### **Function**

Shows chemotactic activity for monocytes, resting T- lymphocytes, and neutrophils, but not for activated lymphocytes. Inhibits proliferation of myeloid progenitor cells in colony formation assays. This protein can bind heparin. Binds CCR1. CCL23(19-99), CCL23(22-99), CCL23(27-99), CCL23(30-99) are more potent chemoattractants than CCL23.

### **Cellular Location**

Secreted.

### **Tissue Location**

High levels in adult lung, liver, skeletal muscle and pancreas. Moderate levels in fetal liver, adult bone marrow and placenta. The short form is the major species and the longer form was detected only in very low abundance. CCL23(19-99), CCL23(22-99), CCL23(27-99), CCL23(30-99) are found in high levels in synovial fluids from rheumatoid patients.

## **MPIF-1/CCL23 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **MPIF-1/CCL23 - Images**