

MIP-3 α /CCL20
Catalog # PVGS1452**Specification**

MIP-3 α /CCL20 - Product Information

Primary Accession [P78556](#)
Species
Human

Sequence
Ala27-Met96

Purity
> 95% as analyzed by SDS-PAGE

Endotoxin Level
< 0.2 EU/ μ g of protein by gel clotting method

Biological Activity
The EC₅₀ value of human MIP-3 α /CCL20 on Ca²⁺ mobilization assay in CHO-K1/G15/hCCR6 cells (human Ga15 and human CCR6 stably expressed in CHO-K1 cells) is less than 200.0 ng/ml.

Expression System
E. coli

Formulation **Lyophilized after extensive dialysis against 53 mM Na₂HPO₄, 147 mM NaH₂PO₄, 300 mM NaCl, pH 6.5.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or PBS up to 100 μ g/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

MIP-3 α /CCL20 - Additional Information

Gene ID 6364

Other Names
C-C motif chemokine 20, Beta-chemokine exodus-1, CC chemokine LARC, Liver and activation-regulated chemokine, Macrophage inflammatory protein 3 α , MIP-3- α , Small-inducible cytokine A20, CCL20(1-67), CCL20(1-64), CCL20(2-70), CCL20, LARC, MIP3A, SCYA20

Target Background

Macrophage Inflammatory Protein-3 (MIP-3 α), also known as chemokine (C-C motif) ligand 20 (CCL20) or liver activation regulated chemokine (LARC), is a small cytokine belonging to the CC chemokine family. MIP-3 α is expressed in the liver, lymph nodes, appendix, PBL and lung and can signal through the CCR6 receptor. It is strongly chemotactic for lymphocytes and weakly attracts neutrophils. MIP-3 α is implicated in the formation and function of mucosal lymphoid tissues via chemoattraction of lymphocytes and dendritic cells toward the epithelial cells surrounding these tissues. Additionally, it promotes the adhesion of memory CD4⁺ T cells and inhibits colony formation of bone marrow myeloid immature progenitors.

MIP-3 α /CCL20 - Protein Information

Name CCL20

Synonyms LARC, MIP3A, SCYA20

Function

Acts as a ligand for C-C chemokine receptor CCR6. Signals through binding and activation of CCR6 and induces a strong chemotactic response and mobilization of intracellular calcium ions (PubMed: [11035086](http://www.uniprot.org/citations/11035086), PubMed: [11352563](http://www.uniprot.org/citations/11352563), PubMed: [20068036](http://www.uniprot.org/citations/20068036)). The ligand- receptor pair CCL20-CCR6 is responsible for the chemotaxis of dendritic cells (DC), effector/memory T-cells and B-cells and plays an important role at skin and mucosal surfaces under homeostatic and inflammatory conditions, as well as in pathology, including cancer and various autoimmune diseases (PubMed: [21376174](http://www.uniprot.org/citations/21376174)). CCL20 acts as a chemotactic factor that attracts lymphocytes and, slightly, neutrophils, but not monocytes (PubMed: [11352563](http://www.uniprot.org/citations/11352563), PubMed: [9038201](http://www.uniprot.org/citations/9038201)). Involved in the recruitment of both the pro-inflammatory IL17 producing helper T-cells (Th17) and the regulatory T-cells (Treg) to sites of inflammation. Required for optimal migration of thymic natural regulatory T cells (nTregs) and DN1 early thymocyte progenitor cells (By similarity). C- terminal processed forms have been shown to be equally chemotactically active for leukocytes (PubMed: [11035086](http://www.uniprot.org/citations/11035086)). Positively regulates sperm motility and chemotaxis via its binding to CCR6 which triggers Ca²⁺ mobilization in the sperm which is important for its motility (PubMed: [23765988](http://www.uniprot.org/citations/23765988), PubMed: [25122636](http://www.uniprot.org/citations/25122636)). Inhibits proliferation of myeloid progenitors in colony formation assays (PubMed: [9129037](http://www.uniprot.org/citations/9129037)). May be involved in formation and function of the mucosal lymphoid tissues by attracting lymphocytes and dendritic cells towards epithelial cells (By similarity). Possesses antibacterial activity towards E.coli ATCC 25922 and S.aureus ATCC 29213 (PubMed: [12149255](http://www.uniprot.org/citations/12149255)).

Cellular Location

Secreted.

Tissue Location

Expressed in the seminal plasma, endometrial fluid and follicular fluid (at protein level). Expressed predominantly in the liver, lymph nodes, appendix, peripheral blood lymphocytes, and fetal lung. Low levels seen in thymus, prostate, testis, small intestine and colon.

MIP-3 α /CCL20 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MIP-3 α /CCL20 - Images