

FGF-10

Catalog # PVGS1475

Specification

FGF-10 - Product Information

Primary Accession Species Human

Sequence Leu40-Ser208

Purity > 95% as analyzed by SDS-PAGE

Endotoxin Level < 0.2 EU/ μg of protein by gel clotting method

Biological Activity

ED < sub > 50 < /sub > < 20.0 ng/ml, measured by a cell proliferation assay using 4MBr-5 cells, corresponding to a specific activity of $> 5.0 \times 10 < sup > 4 < /sup > units/mg$.

015520

Expression System E. coli

Formulation

Lyophilized after extensive dialysis against PBS.

Reconstitution

It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH_2O up to $100 \mu g/ml$.

Storage & Stability

Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

FGF-10 - Additional Information

Gene ID 2255

Other Names Fibroblast growth factor 10, FGF-10, Keratinocyte growth factor 2, FGF10

Target Background

Fibroblast Growth Factor-10 (FGF-10) is a mitogen mainly produced by mesenchymal stem cells in the lung. FGF-10 belongs to the heparin binding FGF family, and is also known as Keratinocyte Growth Factor-2 (KGF-2). It shares homology with KGF and receptor binding to FGFR2-IIIb. However, while KGF induces proliferation and differentiation of various epithelial cells, FGF-10



promotes budding and branching morphogenesis during the multi-organ development via mesenchymal-epithelial cell interactions. FGF-10 is critical for lung and limb development, and is regulated by Shh during early development.

FGF-10 - Protein Information

Name FGF10

Function

Plays an important role in the regulation of embryonic development, cell proliferation and cell differentiation. Required for normal branching morphogenesis. May play a role in wound healing.

Cellular Location Secreted.

FGF-10 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

FGF-10 - Images