

IL-1 β
Catalog # PVGS1479**Specification**

IL-1 β - Product Information

Primary Accession [P26889](#)
Species
Porcine

Sequence
Ala115-Pro267, expressed with an N-terminal Met

Purity
> 95% as analyzed by SDS-PAGE

Endotoxin Level
< 0.2 EU/ μ g of protein by gel clotting method

Biological Activity
ED₅₀ < 1.0 ng/ml, measured by the dose-dependent stimulation of mouse D10S cells, corresponding to a specific activity of 1.0×10^6 IU/mg.

Expression System
E. coli

Formulation **Lyophilized after extensive dialysis against PBS.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or PBS up to 100 μ g/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70°C. Upon reconstitution, the product should be stable for up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.

IL-1 β - Additional Information

Gene ID 397122

Other Names
Interleukin-1 beta, IL-1 beta, IL1B

Target Background
Interleukin-1 beta (rpIL-1 β) is a proinflammatory cytokine produced in a variety of cells including monocytes, tissue macrophages, keratinocytes and other epithelial cells. IL-1 α and IL-1 β are structurally related polypeptides that share approximately 27% amino acid (aa) identity in porcine. While IL-1 α and IL-1 β are regulated independently, they bind to the same receptor and exert

identical biological effects. IL-1 β stimulates thymocyte proliferation by inducing il-2 release, b-cell maturation and proliferation, and fibroblast growth factor activity.

IL-1 β - Protein Information

Name IL1B

Function

Potent pro-inflammatory cytokine. Initially discovered as the major endogenous pyrogen, induces prostaglandin synthesis, neutrophil influx and activation, T-cell activation and cytokine production, B- cell activation and antibody production, and fibroblast proliferation and collagen production. Promotes Th17 differentiation of T-cells. Synergizes with IL12/interleukin-12 to induce IFNG synthesis from T- helper 1 (Th1) cells. Plays a role in angiogenesis by inducing VEGF production synergistically with TNF and IL6. Involved in transduction of inflammation downstream of pyroptosis: its mature form is specifically released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore.

Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:P01584}. Secreted {ECO:0000250|UniProtKB:P01584}. Lysosome {ECO:0000250|UniProtKB:P01584}. Secreted, extracellular exosome {ECO:0000250|UniProtKB:P10749}. Note=The precursor is cytosolic. In response to inflammasome-activating signals, such as ATP for NLRP3 inflammasome or bacterial flagellin for NLRC4 inflammasome, cleaved and secreted. Mature form is secreted and released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore. In contrast, the precursor form is not released, due to the presence of an acidic region that is proteolytically removed by CASP1 during maturation. The secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10. {ECO:0000250|UniProtKB:P01584}

IL-1 β - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IL-1 β - Images