

#### MIF

Catalog # PVGS1656

## Specification

# **MIF - Product Information**

Primary Accession Species Mouse

# <u>P34884</u>

Sequence Met1- Ala115

**Purity** > 96% as analyzed by SDS-PAGE<br>> 96% as analyzed by HPLC

**Endotoxin Level** < 1 EU/ μg of protein by LAL method

Expression System E. coli

**Theoretical Molecular Weight** 12.5 kDa

Formulation

Reconstitution

Lyophilized from a 0.2  $\mu$ m filtered solution in PBS, pH 7.4, 1 mM DTT.

It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml.

#### **Storage & Stability**

Upon receiving, this product remains stable for up to 6 months at -20°C or -70°C. Upon reconstitution, the product should be stable for up to 1 week at 2-8°C or up to 3 months at -20°C. Avoid repeated freeze-thaw cycles.

## **MIF - Additional Information**

Gene ID 17319

**Other Names** 

Macrophage migration inhibitory factor, MIF, 5.3.2.1, Delayed early response protein 6, DER6, Glycosylation-inhibiting factor, GIF, L-dopachrome isomerase, L-dopachrome tautomerase, 5.3.3.12, Phenylpyruvate tautomerase, Mif {ECO:0000303|PubMed:8413654, ECO:0000312|MGI:MGI:96982}

## **Target Background**

Macrophage migration inhibitory factor (MIF or MMIF), also named as glycosylation-inhibiting factor (GIF), L-dopachrome isomerase, or phenylpyruvate tautomerase, is a protein encoded by the MIF



gene. It is released from white blood cells by bacterial antigen stimulation to trigger an acute immune response, or by glucocorticoids to counter-act the inhibitory effects of glucocorticoids on immune system. MIF is a homotrimer of which each subunit contains 115 amino acids. As mentioned above, MIF is involved in the innate immune response to bacterial pathogens and counter-acts the anti-inflammatory activity of glucocorticoids. Furthermore, it also plays a role as mediator in regulating the function of macrophages in host defense and has phenylpyruvate tautomerase and dopachrome tautomerase activity in vitro. Mouse MIF is active on human cells, while human MIF is active on mouse cells. Mouse MIF is 99 %, 84 %, 90 %, and 90 % a.a. identical to rat, porcine, bovine and human MIF, respectively.

# **MIF - Protein Information**

# Name Mif {ECO:0000303|PubMed:8413654, ECO:0000312|MGI:MGI:96982}

## Function

Pro-inflammatory cytokine involved in the innate immune response to bacterial pathogens (By similarity). The expression of MIF at sites of inflammation suggests a role as mediator in regulating the function of macrophages in host defense (By similarity). Counteracts the anti-inflammatory activity of glucocorticoids (By similarity). Has phenylpyruvate tautomerase and dopachrome tautomerase activity (in vitro), but the physiological substrate is not known (PubMed:<a href="http://www.uniprot.org/citations/10933783" target="\_blank">10933783</a>, PubMed:<a href="http://www.uniprot.org/citations/16780921" target="\_blank">10933783</a>, PubMed:<a href="http://www.uniprot.org/citations/16780921" target="\_blank">19188446</a>). It is not clear whether the tautomerase activity has any physiological relevance, and whether it is important for cytokine activity (PubMed:<a href="http://www.uniprot.org/citations/19188446" target="\_blank">10933783" target="\_blank">10933783" target="\_blank">10933783</a>, PubMed:<a href="http://www.uniprot.org/citations/19188446" target="\_blank">10933783" target="\_blank">10933783" target="\_blank">10933783" target="\_blank">10933783</a>, PubMed:<a href="http://www.uniprot.org/citations/19188446" target="\_blank">10933783" target="\_blank">10933783</a>, PubMed:<a href="http://www.uniprot.org/citations/16780921" target="\_blank">10933783" target="\_blank">10933783</a>, PubMed:<a href="http://www.uniprot.org/citations/16780921" target="\_blank">10933783</a>, PubMed:<a href="http://www.uniprot.org/citations/16780921" target="\_blank">10933783</a>, PubMed:<a href="http://www.uniprot.org/citations/16780921" target="\_blank">10933783</a>, PubMed:<a href="http://www.uniprot.org/citations/16780921" target="\_blank">10933783</a>, PubMed:<a href="http://www.uniprot.org/citations/19188446" target="\_blank">10933783</a>.

## **Cellular Location**

Secreted. Cytoplasm {ECO:0000250|UniProtKB:P14174} Note=Does not have a cleavable signal sequence and is secreted via a specialized, non-classical pathway. Secreted by macrophages upon stimulation by bacterial lipopolysaccharide (LPS), or by M.tuberculosis antigens (By similarity). {ECO:0000250|UniProtKB:P14174}

## MIF - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- MIF Images