

## Spike protein RBD

Catalog # PVGS1679

## **Specification**

# **Spike protein RBD - Product Information**

**Primary Accession Species** SARS-CoV-2

P0DTC2

## Seauence

Arg319-Phe541 (G339D, S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H)

### **Purity**

≥ 95% as analyzed by SDS-PAGE

#### **Endotoxin Level**

 $\leq$  1 EU/ µg of protein by LAL method.

## **Biological Activity**

SARS-CoV-2 Spike protein RBD, Omicron Variant, His Tag can bind with human ACE2 in functional ELISA assay.

## **Expression System**

**Human Cells** 

## **Theoretical Molecular Weight**

26.2 kDa

Formulation

Supplied as a 0.2 µm filtered solution of

PBS, pH 7.4.

#### Storage & Stability

Upon receiving, this product remains stable for up to 6 months at -20 °C or below. Avoid repeated freeze-thaw cycles.

# **Spike protein RBD - Additional Information**

Gene ID 43740568

#### **Other Names**

Spike glycoprotein {ECO:0000255|HAMAP-Rule:MF 04099}, S glycoprotein {ECO:0000255|HAMAP-Rule:MF 04099}, E2 {ECO:0000255|HAMAP-Rule:MF 04099}, Peplomer protein {ECO:0000255|HAMAP-Rule:MF\_04099}, Spike protein S1 {ECO:0000255|HAMAP-Rule:MF 04099}, Spike protein S2 {ECO:0000255|HAMAP-Rule:MF 04099}, Spike protein S2' {ECO:0000255|HAMAP-Rule:MF 04099}, S {ECO:0000255|HAMAP-Rule:MF 04099}

## **Target Background**

SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as 2019-nCoV (2019



Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. Recently, the new B.1.1.529 variant was confirmed in South Africa and preliminary evidence suggests an increased risk of reinfection with this variant. The B.1.1.529 variant was first reported to WHO on 24 November 2021 and WHO has designated this variant as a VOC (Variant of Concern), named Omicron. There are more than 30 mutations in the spike protein.

## **Spike protein RBD - Protein Information**

Name S {ECO:0000255|HAMAP-Rule:MF 04099}

#### **Function**


[Spike protein S1]: Attaches the virion to the cell membrane by interacting with host receptor, initiating the infection. The major receptor is host ACE2 (PubMed: <a href="http://www.uniprot.org/citations/32142651" target="\_blank">32142651</a>, PubMed:<a href="http://www.uniprot.org/citations/32155444" target="\_blank">32155444</a>, PubMed:<a href="http://www.uniprot.org/citations/33607086" target="blank">33607086</a>). When S2/S2' has been cleaved, binding to the receptor triggers direct fusion at the cell membrane (PubMed: <a href="http://www.uniprot.org/citations/34561887" target=" blank">34561887</a>). When S2/S2' has not been cleaved, binding to the receptor results in internalization of the virus by endocytosis using host TFRC and GRM2 and leading to fusion of the virion membrane with the host endosomal membrane (PubMed: <a href="http://www.uniprot.org/citations/32075877" target=" blank">32075877</a>, PubMed:<a href="http://www.uniprot.org/citations/32221306" target="blank">32221306</a>, PubMed:<a href="http://www.uniprot.org/citations/34903715" target=" blank">34903715</a>, PubMed:<a href="http://www.uniprot.org/citations/36779763" target=" blank">36779763</a>). Alternatively, may use NRP1/NRP2 (PubMed:<a href="http://www.uniprot.org/citations/33082294" target=" blank">33082294</a>, PubMed:<a href="http://www.uniprot.org/citations/33082293" target="blank">33082293</a>) and integrin as entry receptors (PubMed: <a href="http://www.uniprot.org/citations/35150743" target=" blank">35150743</a>). The use of NRP1/NRP2 receptors may explain the tropism of the virus in human olfactory epithelial cells, which express these molecules at high levels but ACE2 at low levels (PubMed: <a href="http://www.uniprot.org/citations/33082293" target=" blank">33082293</a>). The stalk domain of S contains three hinges, giving the head unexpected orientational freedom (PubMed:<a href="http://www.uniprot.org/citations/32817270" target=" blank">32817270</a>).

## **Cellular Location**

Virion membrane {ECO:0000255|HAMAP-Rule:MF\_04099, ECO:0000269|PubMed:32979942}; Single-pass type I membrane protein {ECO:0000255|HAMAP-Rule:MF\_04099, ECO:0000269|PubMed:34504087}. Host endoplasmic reticulum-Golgi intermediate compartment membrane {ECO:0000255|HAMAP-Rule:MF\_04099, ECO:0000269|PubMed:34504087}; Single-pass type I membrane protein {ECO:0000255|HAMAP-Rule:MF\_04099}. Host cell membrane {ECO:0000255|HAMAP-Rule:MF\_04099}. Note=Accumulates in the endoplasmic reticulum-Golgi intermediate compartment, where it participates in virus particle assembly. Some S oligomers are transported to the host plasma membrane, where they may mediate cell-cell fusion (PubMed:34504087). An average of 26 +/-15 S trimers are found randomly distributed at the surface of the virion (PubMed:32979942) {ECO:0000255|HAMAP-Rule:MF\_04099, ECO:0000269|PubMed:32979942, ECO:0000269|PubMed:34504087}

## Spike protein RBD - Protocols

Provided below are standard protocols that you may find useful for product applications.





- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

**Spike protein RBD - Images**