

TGF- β 1
Catalog # PVGS1688**Specification**

TGF- β 1 - Product Information

Primary Accession [P18341](#)
Species
Bovine

Sequence
Ala279-Ser390

Purity
≥ 95% as analyzed by SDS-PAGE

Endotoxin Level
< 0.2 EU/ μ g of protein by gel clotting method

Biological Activity
ED₅₀ < 0.2 ng/ml, measured in ability to inhibit the mouse IL-4-dependent proliferation of HT-2 cells.

Expression System
CHO

Theoretical Molecular Weight
12.8 kDa (monomer)

Formulation **Lyophilized from a 0.2 μ m filtered solution in 50 mM NaAc, 50 mM NaCl, pH 5.0.**

Reconstitution
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH₂O or 50 mM Citrate up to 100 μ g/ml.

Storage & Stability
Upon receiving, this product remains stable for up to 6 months at lower than -70 °C. Upon reconstitution, the product should be stable for up to 1 week at 4 °C or up to 3 months at -20 °C. For long-term storage after reconstitution, it is recommended that a carrier protein (e.g., 0.1% BSA) be added. Avoid repeated freeze-thaw cycles by making single-use aliquots before the solution is stored at -20 °C.

TGF- β 1 - Additional Information

Gene ID 282089

Other Names
Transforming growth factor beta-1 proprotein, Latency-associated peptide, LAP, Transforming growth factor beta-1, TGF-beta-1, TGFB1

Target Background

TGF- β 1 (transforming growth factor beta 1) is one of three closely related mammalian members of the large TGF- β 1 superfamily that share a characteristic cystine knot structure. TGF- β 1, -2 and -3 are highly pleiotropic cytokines that act as cellular switches to regulate processes such as immune function, proliferation and epithelial-mesenchymal transition. Each TGF- β isoform has some non-redundant function; for TGF- β 1, mice with targeted deletion show defects in hematopoiesis and endothelial differentiation and died of overwhelming inflammation. TGF- β 1 signaling begins with high-affinity binding to a type II ser/thr kinase receptor termed TGF- β RII. This receptor then phosphorylates and activates a second ser/thr kinase receptor, TGF- β RI (also called activin receptor-like kinase (ALK)-5), or alternatively, ALK-1. This complex phosphorylates and activates Smad proteins that regulate transcription.

TGF- β 1 - Protein Information

Name TGFB1

Function

Transforming growth factor beta-1 proprotein: Precursor of the Latency-associated peptide (LAP) and Transforming growth factor beta-1 (TGF-beta-1) chains, which constitute the regulatory and active subunit of TGF-beta-1, respectively.

Cellular Location

[Latency-associated peptide]: Secreted, extracellular space, extracellular matrix
{ECO:0000250|UniProtKB:P01137}

TGF- β 1 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TGF- β 1 - Images